

A study of essential oil from an invasive *Piper aduncum* L.

Kajian minyak atsiri dari tumbuhan invasif *Piper aduncum* L.

I Putu Agus Hendra Wibawa¹, Vienna Saraswaty², Farid Kuswanto¹, Putri Sri Andila¹, Putri Kesuma Wardhani¹,
I Gede Tirta¹, Wawan Sujarwo^{1*}

¹ Bali Botanical Gardens - LIPI, Candikuning, Baturiti, Tabanan Bali 82191

² Development Unit for Clean Technology - LIPI, Jl. Cisitua Sangkuriang Gd. 50 Bandung 40135

*Email: wawan.sujarwo@lipi.go.id

Submitted 26 January 2019; Accepted 20 July 2019

INTISARI

Tumbuhan invasif *Piper aduncum* merupakan spesies tumbuhan yang berpotensi menghasilkan minyak atsiri. Penelitian ini bertujuan untuk menentukan persentase, senyawa kimia yang terkandung, dan sifat anti-jamur dari minyak atsiri yang dihasilkan dari daun dan buah *P. aduncum*. Penelitian menunjukkan bahwa daun dan buah *P. aduncum* mengandung masing-masing 0,30% dan 0,33% minyak atsiri. Apiol adalah senyawa kimia paling melimpah yang diperoleh dalam minyak atsiri daun dan buah *P. aduncum*, masing-masing dengan konsentrasi 57,10% dan 66,31%. Kami menyimpulkan bahwa minyak atsiri yang diperoleh dalam penelitian ini mampu menghambat pertumbuhan *Aspergillus niger* dan *Cladosporium* sp. namun tidak mampu menghambat *Fusarium oxysporum* dan *Fusarium solani*.

Kata kunci: Antijamur, apiol, analisis GC-MS, IAPs, fitokimia, penyakit tanaman

ABSTRACT

An invasive *Piper aduncum* is a plant species that potentially produces essential oil. The study aims to determine the percentage, phytochemical compounds, and anti-fungal properties of essential oil produced from the leaves and fruits of *Piper aduncum*. The study showed that the leaves and fruits of *P. aduncum* contain 0.30% and 0.33% of essential oil, respectively. Apiol is the most abundant phytochemical compound obtained in essential oil of leaves and fruits of *P. aduncum* with 57.10% and 66.31%, respectively. We conclude that essential oil obtained in this study is able to inhibit the growth of both *Aspergillus niger* and *Cladosporium* sp. but not for *Fusarium oxysporum* and *Fusarium solani*.

Keywords: Antifungal, apiol, GC-MS analysis, IAPs, phytochemistry, plant disease

INTRODUCTION

Piper aduncum is firstly introduced to Indonesia as an ornamental plant in 1860, and now has been widely distributed throughout Indonesia (Hartemink, 2010; Tjitrosoedirdjo et al., 2016). The species is neotropic plant that grows to an altitude of 2,000 meters and generally in open and disturbed areas, such as secondary forest (Jan et al., 2002; Tjitrosoedirdjo et al., 2016), for example in Borneo

land clearing for road construction is reported to have contributed to the spread of this species (Padmanaba and Sheil, 2014). Despite being an invasive plant, in Papua New Guinea this species is used for raw material of farming equipment, pesticides, fertilizers, construction of houses, and windbreaks (Siges et al., 2005).

Piper aduncum contains an essential oil which is reportedly able to prevent bites of the *Aedes albopictus* mosquito (Misni et al., 2009). In addition

to being a mosquito repellent, the essential oil is widely used in the drug industry, as food flavor, in cosmetics and fragrances (Rizal et al., 2009). Essential oils are also known to inhibit fungal growth. Essential oil from lemons is known to inhibit the growth of *Aspergillus parasiticum* fungus, *A. carbonarius*, *Cladosporium cladosporioides*, *Eurotium herbariorum* and *Penicillium chrysogenum* (Dimic et al., 2014). Essential oil from *Thymus vulgaris* has also been known to inhibit the growth of the fungus *Rhizopus oryzae* (de Lira Mota et al., 2012).

Efforts to control Invasive Alien Plants (IAPs) are constrained by costs. However, some invasive plant species are considered having an economic value that can be created as part of control costs (Stafford & Blignaut, 2017). This study aims to determine the yield, phytochemical content and potential anti-fungal properties of essential oil obtained from the leaves and fruits of an invasive

Piper aduncum. Also, this present study can hopefully provide a preliminary information to anyone who interested on the topic.

METHODS

Study area

All the processes and antifungal testing of essential oil of *P. aduncum* were conducted at Laboratory of Applied Botany of Bali Botanical Garden, while sequencing of phytochemical was carried out at the Development Unit for Clean Technology - LIPI. The sample materials were the leaves and fruits of *P. Aduncum* obtained from Bali Botanical Gardens with registered number XV.B.229 (Figure 1). We selected leaves which are not too young and not too old, while the fruits were mature (but not ripen yet). We considered picking period for sample materials to yield an optimum amount of essential oil.



Figure 1. *Piper aduncum* (leaves and fruits)

Extraction of Essential Oil

A hundred grams of fresh leaves and fruits of *P. aduncum* were distilled using *Schott Duran* steam distiller. Heating was about five hours at temperature of 100°C. The samples were three

times replication to reduce the data errors. The essential oil was separated with the water using *Schott Duran* separator funnel (Handa, 2008). The separated oil was then analysed using GC-MS. The percentage of essential oil was calculated using the following formula:

$$\text{Essential oil content (\%)} = \frac{\text{Volume of distilled oil (ml)}}{\text{Mass of material that was distilled (gr)}} \times 100$$

Analysis of Chemical Components

The essential oil samples were then analysed using Shimadzu GC-MS QP 2010. About 1 μ L of the sample was injected into Rtx-5MS column (60m \times 0.25mm) with a film thickness of 0.25 μ m, the pre-programmed oven temperature was 50 - 280°C with a temperature rise rate of 5°C/minute. The carrier gas used was helium with a pressure of 101kPa, a total flow rate of 46.5 mL/minute, a column flow rate of 0.85 mL/minute, cleaning flow rate of 3.0 mL/minute, and a split ratio of 1:50.

Antifungal Assay

The essential oil samples were tested for antifungal properties against several species of fungi, such as *Aspergillus niger*, *Cladosporium* sp. *Fusarium solani* and *F. oxysporum* using agar

diffusion method. The method was carried out by laying moistened filter paper with essential oil on the media that had been planted with the fungus with three replications. Clear area formation was observed for three days. The formation of clear area on the surface indicated the presence of fungal death (Pratiwi, 2008).

RESULTS

Extraction of Essential Oil

The distillation of 100 gr of *P. aduncum* fruits produced 0.33 ml of essential oil, which is equivalent to 0.33% of the gross weight of the fruits, while 100 gr of *P. aduncum* leaves produced an average of 0.30 ml of essential oil or 0.30% of the gross weight of the leaves (Table 1).

Table 1. Essential oil content of leaves and fruits of *Piper aduncum*

No.	Samples	Percentage of essential oil (% v/w)			Average
		Replication I	Replication II	Replication III	
1	Leaves	0.329	0.277	0.315	0.307
2	Fruit	0.375	0.350	0.280	0.335

GC-MS analysis

GC-MS analysis of fruits of *P. aduncum* showed that the Apiol was the most dominant compound (66.31%), followed by 3-Cyclohexene-1-one, 2-isopropyl-5-methyl, and Myristicin (8.21% and

4.38% respectively). Apiol was also the highest concentration compound found in the essential oil of *P. aduncum* leaves (57.10%). Other compounds found in quite high concentrations of *P. aduncum* were beta-Cubebene (6.45%), beta-Ocimene (6.30%), and caryophyllene (4.74%) (Table 2).

Table 2. Comparison of GC-MS analysis of essential oil produced from the fruits and leaves of *P. aduncum*

No	Name of Compound	Ret. Time	Fruits	Leaves
			(%)	(%)
1	Bicyclo[3.1.0]hex-2-ene, 2-methyl-5-(1-methylethyl)- <i>or</i> 3-Thujene <i>or</i> .alpha.-Thujene <i>or</i> Origanene <i>or</i> 5-Isopropyl-2-methylbicyclo[3.1.0]hex-2-ene # <i>or</i> 2-Methyl-5-(1-methylethyl)-bicyclo(3.1.0)hex-2-ene \$	7.635	0.58	-
2	(1R)-2,6,6-Trimethylbicyclo[3.1.1]hept-2-ene <i>or</i> 1R-.alpha.-Pinene <i>or</i> Bicyclo[3.1.1]hept-2-ene, 2,6,6-trimethyl-, (1R)- <i>or</i> 2,6,6-Trimethylbicyclo[3.1.1]hept-2-ene-, (1R,5R)- <i>or</i> d-.alpha.-Pinene <i>or</i> 1R-(+)-.alpha.-Pinene <i>or</i> (R)-.alpha.-Pinene	7.856-7.859	1.08	0.66
3	Bicyclo[3.1.1]heptane, 6,6-dimethyl-2-methylene-, (1S)- <i>or</i> 2(10)-Pinene, (1S,5S)-(-)- <i>or</i> (-)-.beta.-Pinene <i>or</i> (-)-2(10)-Pinene <i>or</i> L-.beta.-Pinene <i>or</i> (1S)-(-)-.beta.-Pinene <i>or</i> laevo-.beta.-Pinene <i>or</i> 6,6-Dimethyl-2-methylenebicyclo[3.1.1]heptane-, (S)- <i>or</i> (-)-beta.-Pinene <i>or</i> Bicyclo(3.1.1)heptane, 6,6-dimethyl-2-methylene-, (1S,5S)- <i>or</i> (-)-Pin-2(10)-ene	9.356-9.357	1.08	0.55

Table 2. (cont) Comparison of GC-MS analysis of essential oil produced from the fruits and leaves of *P. aduncum*

No	Name of Compound	Ret. Time	Fruits	Leaves
			(%)	(%)
4	.alpha.-Phellandrene <i>or</i> 1,3-Cyclohexadiene, 2-methyl-5-(1-methylethyl)- <i>or</i> .alpha.-Fellandrene <i>or</i> p-Mentha-1,5-diene <i>or</i> 5-Isopropyl-2-methyl-1,3-cyclohexadiene <i>or</i> 2-methyl-5-(1-methylethyl)-1,3-cyclohexadiene <i>or</i> 4-Isopropyl-1-methyl-1,5-cyclohexadiene <i>or</i> 2-Methyl-5-isopropyl-1,3-cyclohexadiene <i>or</i> alpha-Phellandrene <i>or</i> 5-Isopropyl-2-methyl-cyclohexa-1,3-diene <i>or</i> Menthadiene	10.377-10.380	1.27	0.67
5	1,3-Cyclohexadiene, 1-methyl-4-(1-methylethyl)- <i>or</i> .alpha.-Terpinene <i>or</i> .alpha.-Terpinen <i>or</i> p-Mentha-1,3-diene <i>or</i> Terpilene <i>or</i> 1-Isopropyl-4-methyl-1,3-cyclohexadiene <i>or</i> 1-methyl-4-(1-methylethyl)-1,3-cyclohexadiene <i>or</i> 1,3-Cyclohexadiene, 1-methyl-4-isopropyl- <i>or</i> 1-Methyl-4-isopropylcyclohexadiene-1,3 <i>or</i> .alpha.-Terpine <i>or</i> 1-Methyl-4-isopropyl-1,3-cyclohexadiene <i>or</i> alpha-Terpinene <i>or</i> 1-Isopropyl-4-methyl-cyclohexa-1,3-diene	10.830	1.03	-
6	.alpha.-Pinene <i>or</i> Bicyclo[3.1.1]hept-2-ene, 2,6,6-trimethyl- <i>or</i> 2-Pinene <i>or</i> 2,6,6-Trimethylbicyclo[3.1.1]hept-2-ene <i>or</i> Pinene, .alpha.	11.684	-	2.85
7	Cyclohexene, 1-methyl-5-(1-methylethenyl)-, (R)- <i>or</i> m-Mentha-6,8-diene, (R)-(+)- <i>or</i> (+)-m-Mentha-1(6),8-diene <i>or</i> (+)-Sylvestrene <i>or</i> D-sylvestrene <i>or</i> m-Mentha-1,8-diene, (+)- <i>or</i> Sylvestrene <i>or</i> Sylvestrene, (+)- <i>or</i> 5-Isopropenyl-1-methyl-1-cyclohexene, (R)- <i>or</i> (R)-1-Methyl-5-(1-methylvinyl)cyclohexene	11.290-11.291	2.90	1.99
8	.beta.-Ocimene <i>or</i> 1,3,6-Octatriene, 3,7-dimethyl- <i>or</i> Ocimene <i>or</i> 3,7-Dimethyl-1,3,6-octatriene <i>or</i> beta-Ocimene	12.038-12.093	0.79	6.30
9	.gamma.-Terpinene <i>or</i> 1,4-Cyclohexadiene, 1-methyl-4-(1-methylethyl)- <i>or</i> .gamma.-Terpinen <i>or</i> p-Mentha-1,4-diene <i>or</i> Crithmene <i>or</i> Moslene <i>or</i> 1-methyl-4-(1-methylethyl)-1,4-cyclohexadiene <i>or</i> 1-Methyl-4-isopropyl-1,4-cyclohexadiene <i>or</i> 1-Methyl-4-isopropylcyclohexadiene-1,4 <i>or</i> 1,4-p-Menthadiene <i>or</i> 1,4-Cyclohexadiene, 1-methyl-4-isopropyl- <i>or</i> 4-Isopropyl-1-methyl-1,4-cyclohexadiene <i>or</i> gamma-Terpinene <i>or</i> 1-Isopropyl-4-methyl-1,4-cyclohexadiene # <i>or</i> 1-Isopropyl-4-methyl-cyclohexa-1,4-diene	12.417-12.429	1.98	1.55
10	2-Carene <i>or</i> Bicyclo[4.1.0]hept-2-ene, 3,7,7-trimethyl- <i>or</i> .delta.-2-Carene <i>or</i> (+/-)-2-Carene <i>or</i> 3,7,7-Trimethylbicyclo[4.1.0]hept-2-ene	13.502	0.63	-
11	1,6-Octadien-3-ol, 3,7-dimethyl- <i>or</i> .beta.-Linalool <i>or</i> Linalol <i>or</i> Linalool <i>or</i> Linalyl alcohol <i>or</i> 2,6-Dimethyl-2,7-octadien-6-ol <i>or</i> allo-Ocimenol <i>or</i> 2,6-Dimethyl-2,7-octadiene-6-ol <i>or</i> 2,6-Dimethylocta-2,7-dien-6-ol <i>or</i> 3,7-Dimethyl-1,6-octadien-3-ol <i>or</i> 3,7-Dimethylocta-1,6-dien-3-ol <i>or</i> Linolool <i>or</i> Linanool <i>or</i> 3,7-Dimethyl-octa-1,6-dien-3-ol <i>or</i> dl-3,7-Dimethyl-3-hydroxy-1,6-octadiene <i>or</i> Linalool ex bois de rose oil <i>or</i> Linalool ex ho oil <i>or</i> Linalool ex orange oil <i>or</i> Phantol <i>or</i> Linalool, .beta.	13.977	0.63	-
12	3-Cyclohexen-1-ol, 4-methyl-1-(1-methylethyl)-, (R)- <i>or</i> p-Menth-1-en-4-ol, (R)-(-)- <i>or</i> (-)-Terpinen-4-ol <i>or</i> (-)-4-Terpineol <i>or</i> L-terpinen-4-ol <i>or</i> L-4-terpineneol <i>or</i> L-4-terpineol	16.826-16.819	1.90	0.57
13	3-Cyclohexen-1-one, 2-isopropyl-5-methyl- <i>or</i> 2-Isopropyl-5-methyl-3-cyclohexen-1-one	19.615-19.668	8.21	2.89
14	.alfa.-Copaene	23.686-23.692	0.60	0.55
15	Caryophyllene <i>or</i> Bicyclo[7.2.0]undec-4-ene, 4,11,11-trimethyl-8-methylene-, [1R-(1R*,4E,9S*)]- <i>or</i> Bicyclo[7.2.0]undec-4-ene, 4,11,11-trimethyl-8-methylene-, (E)-(1R,9S)-(-) <i>or</i> .beta.-Caryophyllen <i>or</i> .beta.-Caryophyllene <i>or</i> trans-Caryophyllene <i>or</i> L-Caryophyllene <i>or</i> Bicyclo(7.2.0)undec-4-ene, 8-methylene-4,11,11-trimethyl-, (E)-(1R,9S)-(-) <i>or</i> 8-Methylene-4,11,11-(trimethyl)bicyclo(7.2.0)undec-4-ene, (1R,4E,9S)- <i>or</i> beta-Caryophyllene <i>or</i> .beta.-(E)-Caryophyllene <i>or</i> .beta.-trans-Caryophyllene <i>or</i> Caryophyllen	25.128-25.157	2.24	4.74
16	1,4,7,-Cycloundecatriene, 1,5,9,9-tetramethyl-, Z,Z,Z- <i>or</i> 1,5,9,9-Tetramethyl-1,4,7-cycloundecatriene	26.209-26.226	1.39	2.35
17	.beta.-copaene	27.075	1.40	-

Table 2. (cont) Comparison of GC-MS analysis of essential oil produced from the fruits and leaves of *P. aduncum*

No	Name of Compound	Ret. Time	Fruits	Leaves
			(%)	(%)
18	1H-Cyclopenta[1,3]cyclopropa[1,2]benzene, octahydro-7-methyl-3-methylene-4-(1-methylethyl)-, [3aS-(3a.alpha.,3b.beta.,4.beta.,7.alpha.,7aR*)]- <i>or</i> 1H-Cyclopenta[1,3]cyclopropa[1,2]benzene, 2,3,3a.alpha.,3b.alpha.,4,5,6,7-octahydro-4.alpha.-isopropyl-7.beta.-methyl-3-methylene- <i>or</i> .beta.-Cubebene <i>or</i> .beta.-Cuvebene <i>or</i> 1H-Cyclopenta[1,3]cyclopropa[1,2]benzene, octahydro-7-methyl-3-methylene-4-(1-methylethyl)-, (3a.alpha.,3b.alpha.,4.alpha.,7.beta.,7aR*)-	27.125	-	6.45
19	Cyclohexane, 1-ethenyl-1-methyl-2-(1-methylethenyl)-4-(1-methylethylidene)- <i>or</i> o-Menth-8-ene, 4-isopropylidene-1-vinyl- <i>or</i> Elixene <i>or</i> 2-Isopropenyl-1-methyl-4-(1-methylethylidene)-1-vinylcyclohexane	27.584	-	2.73
20	.alpha.-Farnesene <i>or</i> 1,3,6,10-Dodecatetraene, 3,7,11-trimethyl-, (E,E)- <i>or</i> Farnesene <i>or</i> 2,6,10-Trimethyl-2,6,9,11-dodecatetraene, trans- <i>or</i> 3,7,11-Trimethyl-1,3,6,10-dodecatetraene, (trans,trans)-	27.867	-	1.84
21	Cubedol	28.148-28.168	0.55	1.11
22	1,3-Benzodioxole, 4-methoxy-6-(2-propenyl)- <i>or</i> Benzene, 5-allyl-1-methoxy-2,3-(methylenedioxy)- <i>or</i> Myristicin <i>or</i> 5-Allyl-1-methoxy-2,3-(methylenedioxy)benzene <i>or</i> 6-Allyl-4-methoxy-1,3-benzodioxole <i>or</i> Myristicine	28.366-28.394	4.38	1.64
23	1,5-Cyclodecadiene, 1,5-dimethyl-8-(1-methylethylidene)-, (E,E)- <i>or</i> Germacrene B <i>or</i> Germacra-1(10),4,7(11)-triene <i>or</i> Germacra-1(10),4,7(11)-triene, (E,E)- <i>or</i> 1,5-Dimethyl-8-(1-methylethylidene)-1,5-cyclodecadiene	29.435	-	0.90
24	Viridiflorol	30.573	-	2.56
25	Apiol <i>or</i> 1,3-Benzodioxole, 4,7-dimethoxy-5-(2-propenyl)- <i>or</i> Benzene, 1-allyl-2,5-dimethoxy-3,4-(methylenedioxy)- <i>or</i> Apiole <i>or</i> Apiole <i>or</i> Parsley apiole <i>or</i> Parsley camphor <i>or</i> 1-Allyl-2,5-dimethoxy-3,4-(methylenedioxy)benzene <i>or</i> 4,7-Dimethoxy-5-(2-propenyl)-1,3-benzodioxole <i>or</i> 5-Allyl-4,7-dimethoxy-1,3-benzodioxole # <i>or</i> 1,3-Benzodioxole, 4,7-dimethoxy-5-(2-propen-1-yl)-	32.020-32.064	66.31	57.10
26	Benzene, 1,2,3-trimethoxy-5-(2-propenyl)- <i>or</i> Benzene, 5-allyl-1,2,3-trimethoxy- <i>or</i> Elemicin <i>or</i> 3,4,5-Trimethoxyallylbenzene <i>or</i> 5-Allyl-1,2,3-trimethoxybenzene <i>or</i> 1,2,3-Trimethoxy-5-(2-propenyl)-benzene <i>or</i> 1,2,3-Trimethoxy-5-allylbenzene <i>or</i> 4-Allyl-1,2,6-trimethoxybenzene <i>or</i> Benzene, 5-(2-propenyl)-1,2,3-trimethoxy <i>or</i> Elemicine	32.347	1.05	-

Antifungal assay

Inhibition test on fungi showed an inhibition zone on the inoculated medium of *Aspergillus niger* and *Cladosporium* sp., but not on the inoculated

medium of *Fusarium oxysporum* and *F. Solani* (Table 3). The formation of inhibition zones (Figure 2) on the inoculated medium of *A. niger* and *Cladosporium* sp. showed that essential oil of *P. aduncum* inhibits the fungus growth.

Table 3. Response of *P. aduncum* essential oil on antifungal assay

Essential oil	Anti-fungal activity towards			
	<i>A. niger</i>	<i>Cladosporium</i> sp.	<i>F. oxysporum</i>	<i>F. solani</i>
<i>P. aduncum</i> Fruit	+	+	-	-
<i>P. aduncum</i> Leaves	+	+	-	-




Figure 2. Observation results of antifungal assay (a) *A. niger*; (b) *Cladosporium* sp.; (c) *F. Oxysporum*; and (d) *F. solani*

DISCUSSION

Apiol and Myristicin found in the essential oil of *P. aduncum* fruits and leaves are toxic to insects (Lichtenstein & Casida, 1963; Lichtenstein et al., 1974; Fuhremann and Lichtenstein, 1979; Bernard et al., 1995; Walia et al., 2004), thus the presence of these compounds in high concentrations promotes

an opportunity to use *P. aduncum* essential oil as a natural insect repellent. In addition to Apiol, in a large percentage, the essential oil of *P. aduncum* leaves also contains caryophyllene. This compound is often found in essential oils of various species of *Piperaceae* family member, including *P. cernuum* and *P. regnellii* (Costantin et al., 2001; da Silva et al., 2017).




Figure 3. Comparison of some dominant chemical compounds in essential oils, obtained in fruits and leaves of *P. aduncum*.

Some dominant compounds obtained in the fruits are also found in the leaves with different concentrations. The ratio of such compounds can be seen in Figure 3. This present study showed that the

leaves and fruits of *P. aduncum* are both dominated by terpenoids, these compounds are known having many biological activities, such as antimicrobial, antifungal and antibacterial (Cavaleiro, 2015).

The inhibitory ability is likely to correlate with the presence of beta-Ocimene compounds. Cavaleiro et al. (2015) reported beta-Ocimene is a compound with antifungal properties. The compound is also reported having anti-inflammatory and antiviral properties (Terpene, 2018). On the contrary, ineffectiveness in inhibiting the growth of *F. oxysporum* and *F. solani* was probably because the nature of active ingredients has no wide spectrum, it means the essential oil only inhibits the growth of some fungus species (Neu and Gootz, 2001).

Previous studies of *P. aduncum* extraction showed various yields of essential oil. For example, leaves and branches samples of *P. aduncum* from eight different regions of the Amazon produce essential oil ranging from 1.2% to 3.3% (Maia et al., 1998). *Piper aduncum* leaves, native to Papua New Guinea, yields essential oil of 0.35%, while *P. aduncum* leaves from Cuba yields essential oil of 0.96% (Pino et al., 2004; Rali et al., 2007). The difference in yields of essential oil is probably due to different soil macronutrients in each region. Kahkashan et al. (2015) stated that soil types affected the amount of essential oil produced by *Mentha arvensis* L. Suryawati & Murniyanto (2011) showed that essential oil produced by *Zingiber officinale* L. is directly proportional to the availability of macronutrients in the soil. A lack of available groundwater can also result in changing secondary metabolites of plants than in normal conditions (Trisilawati & Pitono, 2012).

CONCLUSION

The present study concluded that essential oil of the leaves and fruits of *P. aduncum* contained Apiol compounds in high concentrations. The essential oil has been confirmed to inhibit the growth of *Aspergillus niger* and *Cladosporium* sp. The study promoted possibilities that essential oil of *P. aduncum* could be developed into industrial production of fungicides and insecticides. However, natural products should respect to cultivation, collection, and used in sustainable manner which conserves both of the plants materials and the environment.

ACKNOWLEDGMENTS

This study is financially supported by Bali Botanical Gardens. The authors would like to thank to I Gede Tirta and all colleagues who helped in the research and providing technical assistances.

REFERENCES

- Bernard, C.B., H.G. Krishnamurthy, D. Chauret, T. Durst, B.J.R. Philogene, P. Sanches-Vindas, C. Hasbaun, I. Poveda, L.S.Roman, J.T. Arnason. 1995. Insecticidal defenses of Piperaceae from the neotropics. *J. Chem. Ecol.* 21: 801-814.
- Cavaleiro, C., L. Salgueiro, M. Gonçalves, K. Hrimpeng, J. Pinto, E. Pinto. 2015. Antifungal activity of the essential oil of *Angelica major* against *Candida*, *Cryptococcus*, *Aspergillus* and dermatophyte species. *Journal of Natural Medicines* 69(2): 241–248.
- Costantin, M.B., P.Sartorelli, R. Limberger, A.T. Henriques, M. Steppe, M.J.P. Ferreira, M.T. Ohara, V.P. Emerenciano and M.J. Kato. 2001. Essential oils from *Piper cernuum* and *Piper regnellii*: Antimicrobial activities and analysis by GC/MS and ¹³C-NMR. *Planta Med.* 67(8): 771-773. doi:10.1055/s-2001-18363
- da Silva, J.K., R. da Trindade, N.S. Alves, P.L. Figueiredo, JGS. Maia and W.N. Setzer. 2017. Essential oils from neotropical Piper species and their biological activities. *Int. J. Mol. Sci.* 18: 2571. doi:10.3390/ijms18122571
- de Lira Mota, K.S., F. de Oliveira Pereira, W.A. de Oliveira, I.O. Lima and E de Oliveira Lima. 2012. Antifungal activity of *Thymus vulgaris* L. essential oil and its constituent phytochemicals against *Rhizopus oryzae*: interaction with ergosterol. *Molecules* 17:14418-14433. doi:10.3390/molecules171214418
- Dimic, G., S.Kocic-Tanackov, L.Mojovic and J. Pejcin. 2014. Antifungal activity of Lemon essential oil, coriander and cinnamon extracts on foodborne molds in direct contact and the vapor phase. *Journal of Food Processing and Preservation* 39(6):1778-1787. doi:10.1111/jfpp.12410.

- Fuhremann, T.W. and E.P. Lichtenstein, E.P. 1979. Insecticide toxicity and degradation in houseflies as affected by naturally occurring food plant components. *Journal of Agricultural and Food Chemistry* 27:87–91.
- Hartemink, A.E. 2010. The invasive shrub *Piper aduncum* in Papua New Guinea: a review. *Journal of Tropical Forest Science* 22(2): 202–213.
- Handa, S.S. 2008. An overview of extraction techniques for medicinal and aromatic plants. In: Handa, S.S., S.P.S. Khanuja, G. Longo, D.D. Rakesh (Eds.). *Extraction technologies for medicinal and aromatic plants*. ICS-UNIDO. Trieste, Italy. pp: 42-47.
- Jan, L., V. Novovotny., L. Cizek, K. Molem., B. Isua., W. Boen, R. Kutil., J. Auga., M. Kasbal., M. Manumbor and S. Hiuk. 2002. Successful invasion of the neotropical species *Piper aduncum* in rain forests in Papua New Guinea. *Applied Vegetation Science* 5: 255-262.
- Kahkashan, P., B. Najat, S. Iram and S. Iffat. 2016. Influence of soil type on the growth parameters, essential oil yield and biochemical contents of *Mentha arvensis* L. *Journal of Essential Oil Bearing Plants* 19(1): 76-81. doi: 10.1080/0972060X.2015.1086285
- Lichtenstein E.P. and J.E.Casida. 1963. Myristicin, an insecticide and synergistic occurring naturally in the edible parts of parsnip. *Journal of Agricultural and Food Chemistry* 11: 410–415.
- Lichtenstein, E.P., T.T. Liang, K.R. Schulz., H.K. Schnoes, and G.T. Carter. 1974. Insecticidal and synergistic components isolated from dill plants. *Journal of Agricultural and Food Chemistry* (22): 658–664.
- Maia, J.G.S., M. G. B. Zohhbi., E. H. A. Andrade., A. S. Santos., M. H. L. da Silva., A. I. R. Luz and C. B. Bastos. 1998. Constituents of the essential oil of *Piper aduncum* L. growing wild in the Amazon region. *Flavour Fragr. J.* (13): 269-272.
- Misni, N., S. Sulaiman., H. Othman and B. Umar. 2009. Repellency of essential oil of *Piper aduncum* against *Aedes albopictus* in the laboratory. *Journal of the American Mosquito Control Association* 25(4):442–447.
- Neu, H.C. and T.D. Gootz. 2001. Antimicrobial chemotherapy. In: Baron, S. (Ed.). *Medical Microbiology*. 5th(Ed.) Galvestone. The University of Texas Medical Branch.
- Padmanaba, M. and D. Sheil. 2014. Spread of the invasive alien species *Piper aduncum* via logging roads in Borneo. *Tropical Conservation Science* 7 (1):35-44.
- Pino, J.A., R. Marbot, A. Bello and A. Urquiola. 2004. Essential Oils of *Piper peltata* L. Miq. and *Piper aduncum* L. from Cuba. *J. Essent. Oil Res.* 16:124-126.
- Pratiwi, S.T. 2008. *Mikrobiologi Farmasi*. Yogyakarta: Penerbit Erlangga. 176 hal.
- Rali, T., S. W. Wossa, D. N. Leach and P. G. Waterman. 2007. Volatile chemical constituents of *Piper aduncum* L and *Piper gibbilimum* C. DC (Piperaceae) from Papua New Guinea. *Molecules* 12: 389-394.
- Rizal, M., M.S.Rusli, dan A. Mulyadi. 2009. *Minyak Atsiri Indonesia*. IPB. Bogor.
- Stafford, W. and J. Blignaut. 2017. Reducing landscape restoration costs: Feasibility of generating electricity from invasive alien plant biomass on the Agulhas Plain, South Africa. *Ecosystem Services* 27: 224–231. doi:10.1016/j.ecoser.2017.04.008
- Siges, T. H., A. E. Hartemink., P. Hebinck and J. Allen. 2005. The Invasive shrub *Piper aduncum* and rural livelihoods in the Finschhafen area of Papua New Guinea. *Human Ecology* 33(6).
- Suryawati S. dan E. Murniyanto. 2011. Hubungan sifat tanah Madura dengan kandungan minyak atsiri dan tingkat kelarutannya pada jahe (*Zingiber officinale* L). *Agrovigor* 4 (2) : 99 - 104.
- Terpene. 2018. *Everything Terpenes*. <http://terpene.info/terpene/ocimene/>. (Diakses Maret 2016).
- Tjitrosoedirdjo, S.S., I. Mawardi and S. Tjitrosoedirdjo. 2016. 75 Important invasive

alien plant species in Indonesia. SEAMEO BIOTROP Southeast Asian Regional Centre for Tropical Biology. p. 55.

Trisilawati O. dan J. Pitono. 2012. Pengaruh cekaman defisit air terhadap pembentukan bahan aktif pada purwoceng. *Bul Littro*. 23(1):34-47.

Walia, S., S. Saha, B.S. Parmar. 2004. Liquid chromatographic method for the analysis of two plant based insecticide synergist dillapiole and dihydrodillapiole. *J. Chromatogr.* 1047(2): 229-233.