

**THE EFFECT OF INCREASING LEVEL OF PALM KERNEL MEAL AND
COPRA MEAL ON DIET TO THE MICROBIAL PROTEIN PRODUCTION IN
THE RUMEN OF STEERS FED LOW QUALITY FORAGE**

MARSETYO

*Department of Animal Science, The University of Tadulako, Palu, Central Sulawesi,
Indonesia*

SUMMARY

The effect of increasing the level of palm kernel meal (PKM) and copra meal (CM) supplementation on the diet to the microbial crude protein (MCP) production, efficiency of microbial protein production (eMCP) in the rumen and rumen fluid conditions of Brahman crossbred steers given low quality forage based diet has been studied. Ten steers (243 ± 6.5 kg) were allocated to two supplement types diets (PKM and CM). The experimental design included two incomplete 5×5 Latin Squares, each with three runs, for three months. Each run consisted of a 14 d adaptation and 7 d collection period. Steers received Green panic grass hay *ad libitum* with one of five levels of PKM or CM (0.00, 0.25, 0.50, 0.75 and 1.00% of body weight (W) per day (d)). MCP production responded linearly ($P < 0.05$) with increasing PKM or CM intakes. eMCP and rumen $\text{NH}_3\text{-N}$ concentration (taken at 3 and 24 h after feeding) responded quadratically ($P < 0.05$) with increasing PKM or CM intakes. It was concluded that supplementation of both PKM and CM up to level of 1.0% W/day resulted in significant increase in MCP production and eMCP in the rumen. These increases were partly due to the increasing of the concentration of rumen $\text{NH}_3\text{-N}$ in the rumen as a result of increasing supplement intakes.

Key word : Supplement, microbes, rumen, steers

**PENGARUH PENINGKATAN BUNGKIL KELAPA SAWIT DAN BUNGKIL
KELAPA PADA PAKAN TERHADAP PRODUKSI PROTEIN MIKROBA DI
DALAM RUMEN SAPI JANTAN KEBIRIAN YANG MENDAPAT HIJAUAN
BERKUALITAS RENDAH**

RINGKASAN

Pengaruh peningkatan bungkil kelapa sawit (BKS) dan bungkil kelapa (BK) sebagai pakan suplemen terhadap produksi protein mikroba, efisiensi produksi protein mikroba di dalam rumen, dan kondisi cairan rumen pada sapi Brahman jantan kebirian persilangan yang mendapat hijauan berkualitas rendah telah diteliti selama tiga bulan. Sebanyak 10 pedet jantan ($243 \pm 6,5$ kg) dialokasikan pada dua jenis suplemen berupa BKS dan BK. Rancangan percobaan yang digunakan adalah rancangan bujur sangkar ganda latin tidak lengkap (5×5), yang masing-masing terdiri atas tiga periode. Setiap periode terdiri atas 14 hari sebagai masa adaptasi dan 7 hari masa koleksi data. Pada setiap periode, sapi jantan menerima rumput Benggala (*Panicum maximum*) dalam bentuk hay secara *ad libitum* dan salah satu dari lima aras BKS atau BK (0,00; 0,25; 0,50; 0,75 dan 1,00% dari bobot badan (BB) per hari (h)). Peningkatan konsumsi BKS atau BK dapat

meningkatkan secara linier ($P < 0,01$) produksi protein mikroba di dalam rumen. Peningkatan konsumsi BKS atau BK meningkatkan secara kuadrat ($P < 0,05$) efisiensi produksi protein mikroba di dalam rumen dan konsentrasi $\text{NH}_3\text{-N}$ di dalam rumen (3 dan 24 jam setelah ternak mengkonsumsi suplemen). Sebaliknya, pH rumen tidak secara nyata ($P > 0,05$) dipengaruhi oleh peningkatan konsumsi BKS atau BK. Dapat disimpulkan bahwa suplementasi BKS atau BK sampai pada level 1,0% BB/hari pada pedet yang mengkonsumsi rumput berkualitas rendah dapat meningkatkan produksi protein mikroba dan efisiensi produksi protein mikroba di dalam rumen. Peningkatan tersebut sebagian disebabkan oleh peningkatan konsentrasi $\text{NH}_3\text{-N}$ cairan rumen sebagai akibat dari peningkatan konsumsi suplemen.

Kata kunci : Suplemen, mikroba, rumen, pedet

INTRODUCTION

The productivity of beef cattle and other ruminants fed solely low quality forage often very low due to low actual digestibility of feed consumed. The high level of lignification of feed and deficiencies of critical nutrients to support an efficient population of microorganisms in the rumen are the main reasons behind the low utilisation of feed. Hence, supplementary feeding for ruminants given low quality forage is essential to stimulate microbial growth in the rumen, and also nutrient intake by the animal (Orskov, 1999).

Palm kernel meal (PKM) and copra meal (CM), are common supplements for beef cattle and other ruminants which contain lipid and protein. Both PKM and CM are by-products of oil industries which is high in cell-wall constituents and considered as an energy or protein source for ruminants (Devendra, 1988). These feedstuffs have been widely used for ruminant feeding systems in Malaysia (Devendra, 1988), Thailand and Indonesia (Setthapukdee *et al.*, 1991) and in India (Lakshmi and Krisna, 1995). The main limitation of either PKM or CM as a livestock feed has been attributed to their high content of oil which may have negative effect to microbial activities in the rumen (Moore *et al.*, 1986; McLennan *et al.*, 1998). In fact, there is little information on the effect of PKM and CM on microbial protein production in the rumen of beef cattle fed

low quality forage. This study was addressed to examine the effect of increasing either PKM or CM intake on microbial protein production and its efficiency in the rumen of steers received low quality forage.

MATERIALS AND METHODS

Ten Brahman crossbred steers approximately 243 ± 6.5 (SE) kg in initial bodyweight and 15 months of age were allocated to two feed type treatment groups (PKM and CM) by stratified randomisation on the basis of unfasted liveweight. The experimental design was 2 incomplete 5x5 Latin Squares, one with each feed type (PKM and CM). Within each feed type there were five levels of supplement feeding equivalent to 0, 0.25, 0.50, 0.75 and 1.0 % body weight per day (W/d). The basal diet was Green panic hay which was given *ad libitum*. The studies involved 3 runs, with one replicate (steer) per treatment level per run. Thus there were 3 replications of each level of feeding for each feed type overall (3 steers). After each run, steers were randomly allocated to a different level of feeding but steers were kept on the same supplement type. In each run, the steers were fed their diet in individual pens over a 14 d preliminary period and then transferred to individual metabolism crates for 7 d collection period. The steers were weighed at the beginning of the preliminary and collection period to adjust the supplement allocation. Supplements offered were fed based on dry matter content.

The hay was given twice a day in separate portions which were fed at 0800 h and 1200 h. The amount of hay offered each day was set at 10-15% more than that consumed by the steer on the previous day. The supplements were given once daily at

0730 h and fed separately to the basal diet. Fresh drinking water was freely available at all times.

Table 1. Chemical composition of feed ingredients used in the experiment

Component	Green panic hay	Palm kernel meal	Copra meal
Organic matter (%)	91.4	96.8	94.0
Crude protein (%)	5.7	17.1	24.2
Neutral detergent fibre (%)	71.0	67.4	54.9
Ether extract (%)	1.5	10.3	7.3

Measurements

Urine sampling for predicting microbial protein production

For predicting microbial protein production, the urine output of individual steer was measured per day by total collection into trays covered with a filter. The pH of the urine was maintained below 3.0 by adding 10% H₂SO₄ into individual trays at the start of each daily collection. The urine collected over a 24 h period for each steer was mixed and a 5% aliquot was taken and bulked over the collection period into a plastic container stored in a refrigerator. At the end of each collection period, a 5 mL sub-sample was taken from the bulked samples from each steer, diluted to 50 mL with ammonium phosphate stock buffer and frozen awaiting purine derivative (PD) analysis. The concentration of PD in sub-samples of the urine was determined using High Performance Liquid Chromatography (HPLC) procedures (Bolam, 1998)

The exogenous purine supply (X, mmol/day) attributable to the microbial population of the rumen was estimated as the total purine excretion (Y, mmol/d) less the endogenous contribution to this total, divided by a recovery factor. Bowen (2003)

suggested an endogenous purine contribution of 0.190 mmol/kg $W^{0.75}$ for *Bos indicus* cattle and a recovery coefficient of 0.85 for absorbed purines. The calculation thus becomes:

$$Y = 0.85 X + 0.190 W^{0.75}.$$

The value of X (endogenous purine supply) was then used in determining estimated microbial nitrogen production (EMNP, g/d) through the following equation :

$$EMNP = (70X) / (0.83 \times 0.116 \times 1000),$$

where 0.83 is the assumed digestibility of the microbial protein and 0.116 represents the ratio of purine nitrogen to total microbial nitrogen (Chen *et al.*, 1992). A factor of 6.25 is applied to convert the EMNP to a microbial crude protein (MCP) production (g/d).

pH and rumen ammonia-nitrogen concentration in the rumen fluid

Rumen fluid samples were taken 3 h after supplement feeding on day 7 of the collection period and then again just before feeding on the next day (24 h post feeding), by inserting a plastic tube down the oesophagus and into the rumen and withdrawing a sample using a vacuum pump. Ruminal pH was measured on fresh fluid immediately after sampling and a sub-sample (20 mL) of rumen fluid for chemical analysis was drawn immediately into 2 tubes (10 mL capacity), each containing 0.2 mL of concentrated H_2SO_4 , and stored at $-20^{\circ}C$ prior to determination of NH_3-N concentration.

Chemical analysis

Feed samples, refusals and faeces were analysed for dry matter (DM) and organic matter (OM) according to procedure of AOAC (1984) after ground (1 mm

screen). The determination of nitrogen was done by using an automatic total nitrogen analyzer (LECO FP-428). Neutral detergent fibre was determined by procedures of Goering and Van Soest (1970). The ether extract content of the hay and supplements was analysed using a solvent extraction unit (Soxtec HT6, Tecator, Sweden). The concentration of $\text{NH}_3\text{-N}$ in the rumen fluid was measured by a distillation method using a Buchi 321 distillation unit and an automatic titrator. The reagents were 2% boric acid (H_3BO_3) solution, a saturated sodium tetraborate solution (>260 g/L), 0.01M HCL (normality 0.0095) and 25 mL of the boric acid solution.

Statistical analysis

The effects of supplementation were tested and described by fitting general linear models with pen, run, and supplement level as terms using Genstat 6th edition program (Lawes Agricultural Trust, 2002). The difference between control treatment from PKM and CM groups within experiment were also analysed by using Genstat 6th edition. As there were no differences between the controls in both treatment groups tested, a single intercept was used which represented six control steers from two treatment groups.

RESULTS AND DISCUSSION

Microbial protein production

The MCP production and eMCP in the rumen in response to different levels of PKM or CM supplementation are presented in Table 2. The obvious effects of increasing levels of PKM or CM supplementation were to linearly increase the MCP

production. However, increasing intake of both PKM and CM affected quadratically ($P<0.05$) the eMCP in the rumen.

The MCP and eMCP were low for the control steers and increased as the proportion of supplements in the ration increased. The low eMCP of the six control steers (90 g MCP/kg digestible organic matter (DOM)) in this experiment is in agreement with previous studies (Bolam, 1998 and Mullik, 1999). In the study of Mullik (1999) on steers fed Pangola grass hay (CP=8.5%) *ad libitum*, the eMCP was 77 g MCP/kg DOM. These low values are primarily related to rumen degradable protein (RDP) supply where the crude protein (CP)/DOM of the Green panic was 112 g CP/kg DOM. The low availability of RDP, soluble carbohydrate and dilution rate, all contribute to this low eMCP (Leng, 1990). However, with increased intake of either PKM or CM in the current study, MCP production showed a linear response to both supplement types. Furthermore, eMCP responded quadratically to increasing intake of both supplements. Interestingly, at the higher level of CM intake (0.68 to 1.0 %W/d), the value of eMCP reached the range level suggested in the current feeding standards (AFRC, 1992; NRC, 2000) of 130-170 g MCP/kg DOM. There appears to be no literature using similar supplements to compare these values. Enhanced MCP production and eMCP due to the supplementation of PKM or CM is probably due to an increase in RDP supply reflected in a higher rumen $\text{NH}_3\text{-N}$ concentration (Table 2), and readily fermentable energy as seen in higher DM intake (Table 2).

Table 2. Effect of feeding palm kernel meal (PKM) and copra meal (CM) on the microbial protein production (MCP), the efficiency of MCP production (eMCP), total dry matter (DM) intake, rumen parameters of steers received low quality forage.

Items	Supplement intake (%W/d)					Probability	
	0.00	0.25	0.50	0.75	1.00	Linear	Quad
MCP production (g/d)							
PKM	238±10.8	300±18.0	353±12.2	434±33.0	469±20.1	<0.01	0.42
CM		327±16.2	400±19.9	469±35.8	614±33.2	<0.01	0.58
eMCP (g CP/kg DOMI)							
PKM	90±2.9	94±2.2	102±2.4	121±8.2	134±6.1	0.49	0.04
CM		95±2.7	111±2.0	130±3.3	160±2.3	0.14	0.03
Total DM intake (%W/d)							
PKM	1.8±0.1	2.0±0.1	2.2±0.1	2.3±0.2	2.2±0.1	<0.01	<0.01
CM		2.0±0.1	2.1±0.1	2.1±0.2	2.2±0.3	<0.01	<0.01
Rumen NH₃-N –3 h (mg/L)							
PKM	42±0.4	65±1.4	74±1.1	80±0.9	97±1.2	<0.01	<0.01
CM		67±1.1	77±1.9	97±0.8	114±1.6	<0.01	0.04
Rumen NH₃ N –24 h (mg/L)							
PKM	25±0.3	54±0.8	60±0.9	65±1.2	70±0.9	<0.01	<0.01
CM		58±0.8	64±1.1	73±0.9	81±1.2	<0.01	<0.01
Rumen pH-3 h							
PKM	7.1±0.1	7.1±0.0	7.1±0.1	7.2±0.1	7.1±0.3	0.99	0.65
CM		7.1±0.2	7.0±0.2	6.8±0.1	6.8±0.1	0.83	0.96
Rumen pH-24 h							
PKM	6.9±0.2	7.0±0.1	7.0±0.0	7.2±0.3	6.9±0.1	0.97	0.99
CM		6.9±0.3	6.9±0.2	6.7±0.2	6.9±0.2	0.85	0.84

Rumen condition parameters

Rumen NH₃-N concentrations were increased by both PKM and CM supplementation as shown in Table 2. As the proportion of supplements (PKM or CM) in the diet increased, the values of NH₃-N concentration increased quadratically (P<0.01).

A higher ruminal NH₃-N concentrations observed in the current experiment with increasing PKM or CM supplementation agrees with other research (Ehrlich *et al.*, 1990) and reflects the high protein degradation in the rumen. The ruminal NH₃-N concentration without supplementation was low (42 and 25 mg/L, at 3 and 24 h after feeding, respectively) and below the minimal accepted level of 50 mg/L (Satter and

Slyter, 1974). Although the level of $\text{NH}_3\text{-N}$ concentration in the rumen increased with increasing level of supplements, the value is considered to be lower than that the value recommended by Perdok and Leng (1990) (at least 100 mg $\text{NH}_3\text{-N/L}$) to promote higher digestion of fibre in the rumen. The high lipid concentration in the supplements might inhibit access to the site and degradation of protein.

With respect to rumen pH, no significant difference due to diets was observed. The overall mean values for pH in rumen fluid at 3 h after feeding was 7.1 ± 0.1 for PKM supplemented group, and 6.9 ± 0.2 for CM supplemented group. Earlier studies (Stewart, 1977, Russell and Dombrowski, 1980) demonstrated that when ruminal pH is above 6.0, cellulolytic enzymes and cellulolytic bacteria are not inhibited. The range of pH across treatments and sampling times in the current study was 6.6 to 7.2, indicating that pH should not have affected microbial activities in the rumen.

CONCLUSIONS

The MCP production and eMCP in the rumen of steers fed low quality forage was enhanced by supplementation of either PKM or CM up to the level of 1.0%W/d which were resulted from the higher concentration of rumen $\text{NH}_3\text{-N}$ of rumen fluid.

ACKNOWLEDGMENTS

We wish to thank to Cooperative Research Centre (CRC) for cattle and beef quality for financial and technical support that enabled this experiment done. Sincere thanks are also due to Jim Kidd and other staff at the Animal Research Institute,

Department of Primary Industry of Queensland, Australia for professional and technical support.

REFERENCES

- AFRC. 1992. Nutritive requirements of ruminants: protein. *Nutr. Abst. Rev.* **62**:787-835.
- AOAC. 1984. *Official Methods of Analysis, Centennial Edition*. Association of Official Analytical Chemist, Washington, DC.
- Bolam, M. J. 1998. Manipulation of the supply of protein and energy to ruminants consuming tropical forages through supplementation strategies. PhD Thesis. University of Queensland.
- Bowen, M.K. 2003. Efficiency of microbial crude protein production in cattle grazing tropical pasture. PhD Thesis. University of Queensland.
- Chen, X.B., Chen, Y.K., Franklin, M.F., Orskov, E.R. and Shard, W.J. 1992. The effect of feed intake and body weight on purine derivatives excretion and microbial protein supply in sheep. *J. Anim. Sci.* **27**:1709-1715.
- Devendra, C. 1988. Utilisation of feeding stuffs from the oil palm. In: *Proc Feeding Stuffs for Livestock in South East Asia*. FAO/APHCA Publication No.7 FAO Regional officer for Asia and the Pacific, Bangkok. pp.39-49.
- Ehrlich, W.K., Upton, P.C., Cowan, R.T. and Moss, R.J. 1990. Copra meal as a supplement for grazing dairy cows. *Proc. Aust. Soc. Anim. Prod.* **18**:196-199.
- Goering, H. K and Van Soest, P.J. 1970. Forage fibre analyses (apparatus, reagents, procedures and some applications).
- Lakshmi, V.P. and Krishna, N. 1995. Evaluation of complete rations containing varying levels of palm kernel-cake as a replacement for groundnut-cake in sheep. *Indian J. Anim. Sci.* **65**: 1161-1164.
- Lawes Agricultural Trust, 2002. Genstat 6th Edition for Windows. Version 6.1. Numerical Algorithms Group. Oxford.
- Leng, R.A. 1990. Factors affecting the utilisation of poor quality forages by ruminants particularly under tropical conditions. *Nutr. Res. Rev.* **3**:277-303.
- McLennan S.R., Plasto, A.W., Doogan, V.J. and Dillon, R.D. 1998. Whole cottonseed and cottonseed meal supplements for cattle given a hay-based diet. *Proc. Aust. Soc. Anim. Prod.* **22**:111-114.

- Moore, J.A., Swingle, R.S. and Hale, W.H., 1986. Effect of whole cottonseed, cottonseed oil or animal fat on digestibility of wheat straw diets by steers *J. Anim. Sci.* **63**:1267-1273.
- Mullik, M.L. 1999. Strategies to increase efficiency of rumen microbial protein synthesis and productivity of cattle on a tropical grass pasture. PhD Thesis. University of Queensland.
- NRC. 2000. Nutrient Requirements of Beef Cattle. 7th revised edition, National Academy Press. Washington, D.C
- Orskov, E.R. 1999. Supplement strategies for ruminants and management of feeding to maximise utilisation of roughages. *Prev. Vet. Med.* **38**:179-185.
- Perdok, H.B. and Leng, R.A. 1990. Effect of supplementation with protein meal on the growth of cattle given a basal diet of untreated or ammoniated rice straw. *Asian-Aust. J. Anim. Sci.* **3**:369-279.
- Russell, J.B. and Dombrowski, D.B., 1980. Effect of pH on the efficiency of growth by pure cultures of rumen bacteria on continuous culture. *Appl. Environ. Microbiol.* **39**:604.
- Satter, L.D. and Slyter, L.L. 1974. Effect of ammonia concentration on rumen microbial protein production *in vitro*. *Br. J. Nutr.* **32**:199-208.
- Setthapukdee, C., Jalaludin, S. and Mahyuddin, M. 1991. Effect of supplementing grass and urea on the performance and carcass characteristics of lambs fed palm kernel cake-based diet. *Proc. 3rd Int. Symp. Nutr. Herb. Penang, Malaysia.* p.100.
- Stewart, C.S. 1977. Factors affecting the cellulolytic activity of rumen contents. *Appl. Environ. Microbiol.* **33**:497-502.