

PREVALENCE OF HYPERURICEMIA AT NEPALGUNJ MEDICAL COLLEGE, BANKE-NEPAL

Singh, P., Khan, S., and Mittal, R.K.

Department of Biochemistry Nepalgunj Medical College Chisapani Banke-Nepal

Objective: The prevalence of hyperuricemia varied in different populations and it appeared to be increasing in the past decades. Recent studies suggest that hyperuricemia is an independent risk factor for cardiovascular disease. **Method:** One thousand eighty seven patients, attending Nepalgunj medical college and Teaching Hospital, Banke, Nepal from March 2011 to February 2012 were included in this study. The reference range was 3.5 – 7.0 mg/dL in males and 2.6 – 6.0 mg/dL in females. Hyperuricemia was defined as a Serum Uric Acid level of more than 7.0 mg/dL in males and of more than 6.0 mg/dL in females. The uric acid was determined by uricase /PAP method. Test was performed in the central laboratory of Biochemistry, Nepalgunj Medical College & teaching Hospital, Banke, Nepal. **Results:** A number of 1487 studies were selected, the statistical information of which was collected for systematic analysis. The results showed that the high prevalence of hyperuricemia found in females(22.86%) as compare to males(18.98). It was found that 21-40 age group is on high risk for hyperuricemia. **Conclusion:** The prevalence of hyperuricemia is different as the period of age and it increases after 21-40 years in male and in female. Serum uric acid level was high in female as compare to male.

Keywords: Prevalence, Nepalgunj medical college, Hyperuricemia, Serum Uric Acid level

INTRODUCTION

The incidence of gout has been increasing in middle-aged and older people globally during the last two decades and has been related to modernization.¹⁻⁶

Hyperuricemia is considered to be the most significant risk factor for gout and may play a role in the development of many degenerative diseases.^{7,8,9-17} Hyperuricemia may be presented asymptotically, or manifested in gouty arthritis, uric acid nephropathy, and nephrolithiasis.¹⁸ The prevalence of hyperuricemia and gout shown an increasing trend all over the world including in developing countries and the reasons include dietary habits, increasing longevity, The prevalence of hyperuricemia in Nepal and in the developing world is less studied than in the developed world.¹⁹

MATERIALS AND METHOD

It was a retrospective study. We have taken sample of 1487 patients for analyzation of Serum Uric Acid level in the central laboratory of Biochemistry, Nepalgunj Medical College,

Nepalgunj, Nepal from March 2011 to February 2012. The reference range was 3.5 – 7.0 mg/dL in males and 2.6 – 6.0 mg/dL in females. Hyperuricemia was defined as a Serum Uric Acid level of more than 7.0 mg/dL in males and of more than 6.0 mg/dL in females.^{20,21,22,23} The uric acid was determined by uricase/PAP method (Fossati & Prencipe, 1980) using the diagnostic reagent kit manufactured by Crest bio systems, a division of coral clinical systems Goa, India. Patients were categorized into males and females. The instructions, test procedure, reagents and accessories to follow were supplied with the kit.

RESULTS

Total 1487 patients were included in this study. Of them, 42.17% male and 57.83% female were tested for serum uric acid level (Figure 1). Serum uric acid level found elevated in 247 patients and 1240 found within normal range. In 247 elevated cases 119 were male and 128 were female. In 1240 normal range cases 508 were male and 732 were female. Highest elevated level were found in the age group of 21- 40 (Table-I & figure II). The prevalence of total patients was 16.61%. Prevalence of hyperuricemia of total female was 22.86% and the prevalence of hyperuricemia of total male was 18.98% (Table 2). The highest prevalence of hyperuricemia of male found in the age group 21-40 & 41-60 (35.29% & 36.97%) and the

Correspondence: Singh, P.
Address: Department of Biochemistry
Nepalgunj Medical College Chisapani Banke-
Nepal. Phone-009779848354981
Email: [priti186631@gmail.com](mailto:pritti186631@gmail.com)

highest prevalence hyperuricemia of female found in the age group 21-40 was 42.19%. (Table 3 and Figure 3).




Figure 1
Sex Distribution of cases

Table 1
Prevalence of Hyperuricemia according to Age Group Wise Distribution

Age	Serum uric acid		Total	Prevalence rate (%)
	Normal	Elevated		
0-20	152	35	187	18.71
21-40	583	96	679	14.14
41-60	366	80	446	17.94
60+	139	36	175	20.57
Total	1240	247	1487	16.61




Figure 2
Age group wise distribution of elevated level of serum uric acid in total patients

DISCUSSION

The prevalence of hyperuricemia varies in different populations and areas. In Turkey 24, one study reported that 19% of the men and 5.8% of the women had hyperuricemia and the overall prevalence of hyperuricemia was 12.1% in the urban population. In Nepal 18, 3794 people which

were from Chitwan districts were investigated, and the prevalence of hyperuricemia was 21.42%.

Table 2
Sex Wise Prevalence of Hyperuricemia

Sex	Serum uric acid		Total	Prevalence rate (%)
	Normal	Elevated		
Male	508	119	627 (42.17%)	18.98
Female	732	128	860 (57.83%)	22.86
Total	1240	247	1487 (100%)	16.61

Table 3
Prevalence of Hyperuricemia According to Sex wise Distribution in Different Age Groups

Age	Sex	
	Male	Female
0-20	9.24	18.75
21-40	35.29	42.19
41-60	36.97	28.13
60+	18.49	10.94
Total	18.98	22.86




Figure 3
The Comparative Prevalence of Hyperuricemia of Male and Female in Different Age Group.

In Seychelle 20, the cross-sectional health examination survey based on a population random sample which included 1011 subjects aged 25 to 64 years showed that the prevalence of hyperuricemia was 35.2% and 8.7% in men and women, respectively. In Thailand 25, an across-sectional study of 1381 patients who firstly participated in annual health examinations during the period of July 1999 through February 2000 reported that the prevalence of hyperuricemia was 10.6%, but it was 18.4% and 7.8% in men and women, respectively. In Java 26, the prevalence of hyperuricemia was investigated by a survey of a total population of 4683 rural adults and the result was 24.3%. In United States 27, the prevalence rate of

asymptomatic hyperuricemia in the general population was estimated at 2-13%. The prevalence of gout and/or hyperuricemia increased about 2 cases per 1000 enrollees over 10 year (1990-1999) in the overall population. In Japan 28, a total of 9,914 individuals (6,163 men and 3,751 women aged from 18 to 89 years) who were screened at Okinawa General Health Maintenance Association was screened. The result showed that the prevalence of hyperuricemia was 25.8% and it was 34.5%, 11.6% in men and women respectively. In New Zealand 29, hyperuricemia was more common in Maori men (27.1%) than in European men (9.4%) and in Maori women (26.6%) than in European women (10.5%). In Saudi Arabia [73], the prevalence of hyperuricemia was only 8.84%. In Taiwan island of China 30, the prevalence of hyperuricemia was high to 49.4% in Ayatals, but it was only 27.4% in non-aborigines.

In our study the prevalence of hyperuricemia seems to be higher in female as compare to male; it was 22.86% in female and 18.98% in male. The age specific prevalence in this study was also found to be higher (20.57%) in more than 60 year age groups and 18.71% was in age group 0-20 %. From the analysis, it was found that age and sex affected the serum uric acid levels and the prevalence of hyperuricemia.

CONCLUSION

The prevalence of hyperuricemia is different as the period of age and it increases after 21-40 years in male and in female. Serum uric acid level was high in female as compare to male.

REFERENCES

1. Arromdee E, Michet CJ, Crowson CS, O'Fallon WM, Gabriel SE. Epidemiology of gout: is the incidence rising? *J Rheumatol* 2002; 29: 2403-2406.
2. Chen S, Du H, Wang Y, Xu L. The epidemiology study of hyperuricemia and gout in a community population of Huangpu District in Shanghai. *Chin Med J (Engl)* 1998; 111: 228-230.
3. Chen SY, Chen CL, Shen ML, Kamatani N. Trends in the manifestations of gout in Taiwan. *Rheumatology (Oxford)* 2003; 42: 1529-1533.
4. Klemp P, Stansfield SA, Castle B, Robertson MC. Gout is on the increase in New Zealand. *Ann Rheum Dis* 1997; 56: 22-26.
5. Zeng Q, Wang Q, Chen R, Xiao Z, Huang S, Xu J. Primary gout in Shantou: a clinical and epidemiological study. *Chin Med J (Engl)* 2003; 116: 66-69.
6. Wallace KL, Riedel AA, Joseph-Ridge N, Wortmann R. Increasing prevalence of gout and hyperuricemia over 10 years among older adults in a managed care population. *J Rheumatol* 2004; 31: 1582-1587.
7. Terkeltaub RA. Clinical practice. Gout. *N Engl J Med* 2003; 349: 1647-1655.
8. Mahowald ML. Overview of the evaluation and management of gout and hyperuricemia. *Rheumatology & Musculoskeletal Medicine for Primary Care*, Gout 2004/10/8 [cited 2005/5/12].
9. Franse LV, Pahor M, DiBari M, Shorr RI, Wan JY, Somes GW, et al. Serum uric acid, diuretic treatment and risk of cardiovascular events in the Systolic Hypertension in the Elderly Program (SHEP). *J Hypertens* 2000; 18:1149-1154.
10. Lai SW, Tan CK, Ng KC. Epidemiology of hyper-uricemia in the elderly. *Yale J Biol Med* 2001;74:151-157.
11. Lin KC, Tsai ST, Lin HY, Chou P. Different progressions of hyperglycemia and diabetes among hyperuricemic men and women in the Kinmen study. *J Rheumatol* 2004; 31: 1159-1165.
12. Tseng CH. Independent association of uric acid levels with peripheral arterial disease in Taiwanese patients with Type 2 diabetes. *Diabet Med* 2004; 21: 724-729.
13. Langlois M, De Bacquer D, Duprez D, De Buyzere M, Delanghe J, Blaton V. Serum uric acid in hypertensive patients with and without peripheral arterial disease. *Atherosclerosis* 2003; 168: 163-168.
14. Li Y, Stamler J, Xiao Z, Folsom A, Tao S, Zhang H. Serum uric acid and its correlates in Chinese adult populations, urban and rural, of Beijing. The PRC-USA Collaborative Study in Cardiovascular and Cardio-pulmonary Epidemiology. *Int J Epidemiol* 1997; 26: 288-296.
15. Nagahama K, Iseki K, Inoue T, Touma T, Ikemiya Y, Takishita S. Hyperuricemia and cardiovascular risk factor clustering in a screened cohort in Okinawa, Japan. *Hypertens Res Clin Exp* 2004; 27: 227-233.
16. Niskanen LK, Laaksonen DE, Nyssonen K, Alfthan G, Lakka HM, Lakka TA, et al. Uric acid level as a risk factor for cardiovascular and all-cause mortality in middle-aged men: a prospective cohort study. *Arch Intern Med* 2004; 164: 1546-1551.
17. Chu NF, Wang DJ, Liou SH, Shieh SM. Relationship between hyperuricemia and other cardiovascular disease risk factors among adult males in Taiwan. *Eur J Epidemiol* 2000; 16: 13-17.
18. Kumar S, Singh AR, Takhelmayum R, Shrestha P, Sinha JN, et al.: Prevalence of hyperuricemia in Chitwan District of Nepal. *Journal of college of Medical Sciences-Nepal* 2010, 6(2):18-23.
19. Pokhrel K, Yadav BK, Jha B, Parajuli K, Pokharel RK. Estimation of Serum Uric Acid in

- Cases of Hyperuricaemia and Gout J Nepal Med Assoc 2011;51(181):15-20.
20. Conen D, Wietlisbach V, Bovet P, Shamlaye C, Riesen W, Paccaud F, et. al. Prevalence of hyperuricemia and relation of serum uric acid with cardiovascular risk factors in a developing country. BMC Public Health. 2004;25:4-9.
 21. Nan H, Qiao Q, Dong Y, Gao W, Tang B, Qian R, et.al. The prevalence of hyperuricemia in a population of the coastal city of Qingdao, China. J Rheumatol 2006;33:1346-50.
 22. Yu KH, See LC, Huang YC, Yang CH, Sun JH. Dietary factors associated with hyperuricaemia in adults; Semin Arthritis Rheum 2008;37:243-50.
 23. Dehghan A, Köttgen A, Yang Q, Hwang SJ, Kao WL, Rivadeneira F, et al. Association of three genetic loci with uric acid concentration and risk of gout: a genome-wide association study. Lancet 2008;372:1953-61.
 24. Ismail S, Servet A, Betul P, Ali RS, Oguz G, Merih B, et al.: Hyperuricemia and its related factors in an urban population, Izmir, Turkey. Rheumatol Int 2009, 29:869-874.
 25. Vitoon J, Rungroj K, Thananya B, Kamol U, Suthipo U: Prevalence of Hyperuricemia in Thai Patients with Acute Coronary Syndrome.
 26. Darmawan J, Valkenburg HA, Muirden KD, Wigley RD: The epidemiology of gout and hyperuricemia in a rural population of Java. J Rheumatol 1992, 19:1595-1599.
 27. Edward W, Campion MD, Robert J, Glynn SD, Lorraine O, Delabry MA: Asymptomatic hyperuricemia. Risks and consequences in the normative aging study The American Journal of Medicine 1987, 82(3):421-426.
 28. Kazufumi N, Kunitoshi I, Taku I, Takashi T, Yosiharu I, Shuichi T: Hyperuricemia and Cardiovascular Risk Factor Clustering in a Screened Cohort in Okinawa, Japan. Hypertens Res 2004, 27:227-233. PubMed Abstract | Publisher Full Text.
 29. Patrick K, Shelley AS, Benjamin C, Robertson MC: Gout is on the increase in New Zealand. Annals of the Rheumatic Diseases 1997, 56:22-26.
 30. Abdurhman SA-A: Hyperuricemia in Saudi Arabia Rheumatol Int. 2001, 20:61-64.