

EVALUATION OF LAND USE WITH REGIONAL SPATIAL PLAN (RTRW) USING SATELLITE DATA AND GIS IN BATUR UNESCO GLOBAL GEOPARK

I Putu Wira Utama^{1*)}, Takahiro Osawa²⁾, I Wayan Sandi Adnyana³⁾

¹⁾BappedaLitbangof Bangli Regency

²⁾Center for Remote Sensing and Ocean SciencesUdayana University

³⁾Environment Science Postgraduate Program Udayana University

*E-mail: wira_ajus@yahoo.com

ABSTRACT

Development in Batur UNESCO Global Geopark which has an area of 19,422.39 ha has increased significantly in recent years. The existence of land constraints and to determine the suitability of land use, it is necessary to evaluate land use with regional spatial plan (RTRW). Landsat 8 satellite remote sensing data on 27 September 2017 is used to create land use maps. Land use maps obtained through the process of image classification using supervised classification method and verified by ground check. From this technique result 11 classes of land use. Furthermore, to evaluate of land use suitability was conducted by comparing land use with regional spatial plan (RTRW). In this process, there is an overlay between the land use maps with regional spatial plan (RTRW) map using geographic information system (GIS). The results of evaluation land use in Batur UNESCO Global Geopark with regional spatial plan (RTRW) overall has suitable area 10,863.14 ha (55.93%), not suitable area 8,275.58 ha (42.61%) and not detected/cloud interference 283.67 ha (1.46%).

Keywords: land use, evaluation, land use, satellite data, GIS, geopark.

1. INTRODUCTION

The recognition of Batur caldera as the first geopark in Indonesia (year 2012) further adds to the commercial value of Kintamani. This attracts entrepreneurs to start making use of tourist arrivals by providing tourism facilities such as restaurants, guest houses, hot spring pools, agro tourism and other tourism facilities. The development of tourism facilities has increased significantly in recent years, but has not been matched by sustainable and professional management. Entrepreneurs build facilities tourism without refers to spatial plan and obstruct natural panorama which is actually the main attraction Kintamani (Agustini, 2015).

Remote sensing data, especially from satellite remote sensing systems, is very effective in identifying land cover for several reasons. This provides a synoptic and comprehensive view for a large area, which is not possible through land surveying (Richards, 2013). Remotely sensed satellite data have been used to monitor and obtain data of land cover and land use (Adnyana *et al.*, 2006). Landsat satellites have collected and archived moderate resolution data of the entire earth's land surface long enough to be useful for monitoring trends and evaluating how land use has changed. Applications of Landsat data include agriculture, land mapping and change detection to uses in fresh and coastal waters (Knight and Kvaran, 2014).

The GIS can also provide a spatial overview of

detailed land use and closure (Wirosoedarmo *et al.*, 2014). A GIS technology was developed from 1967 by Roger Tomlinson used for scientific investigation, resource management, development planning, cartography and so on, even more profitable than direct field surveys. GIS can be utilized to help incorporate or overlay between different maps and influence the formation of ground mapping units (Hidayati and Toyibullah, 2011). Using Landsat imagery and GIS process, Naningsi (2016) shows the change of land use in Bangli regency years 1997-2014 that the plantation increased 19,486.33 ha (36.89%) and residential increased 1,872.00 ha (3.47%). Land use should refer to the designation of the space within an area which includes the designation of space for the protection function and the designation of space for the cultivation function.

Efforts to balance the utilization of natural resources and the environment is through spatial planning based on the achievement of environmental function preservation (Wirosoedarmo *et al.*, 2014). Preservation of environmental functions can be guaranteed by spatial use activities that pay attention to the carrying capacity of the environment. The carrying capacity of the environment becomes the most important consideration in spatial planning, both in the regional spatial plan (RTRW) and in the evaluation of land use. The objective of this research is to evaluate land use based on spatial plan (RTRW), so it is known the suitability of land use.

2. METHODOLOGY

2.1. Reserch Location

The research location is in Batur UNESCO Global Geopark, Kintamani District, Bangli Regency, Bali Province. Geographically located in 08° 9' 48" S – 08° 21' 26" S and 115° 17' 52" E – 115° 27' 28" E. The total area of Batur UNESCO Global Geopark is 19,422.39 ha or 37.29 percent of Bangli Regency (52,081 ha). The administrative boundaries of Batur UNESCO Global Geopark are northern Buleleng Regency, eastern Karangasem Regency, south Bangli District and western Manikliyu, Belancan, Bantang, Daup, Serai, and Awan Village (Figure 1).




Figure 1.
Research Location

2.2. Materials and Method

The materials of this research are satellite data Landsat 8 OLI 30 x 30 m and 15 x 15 m Pan-resolution, date 27 September 2017 Path/Row=116/66 download from the website www.earthexplorer.usgs.gov; regional spatial plan map of Bangli Regency (RTRW) scale 1: 60:000 from Bappeda Litbang of Bangli Regency; Worldview 2 and Pleiades imagery of Bangli Regency year record 2013 until 2015, resolution 0.5 x 0.5 m, from Geospatial Information Agency (BIG) Bogor.

To conduct land use evaluation of land use with regional spatial plan (RTRW) are processed with GIS according to mechanism as shown in Figure 2.

A. Land Use Map

- 1) Cropping
Cropping of imagery was conducted to determine the locations of the research were observed in accordance with the administrative boundaries Batur UNESCO Global Geopark using ArcGIS 10.5 software.
- 2) Color Composite Image (RGB)
There are three primary colors (red, green and blue) in displaying a color composite image. When




Figure 2.
Data Analysis Mechanism

these three colors are combined in various proportions, they produce different colors in the visible spectrum. Each spectrum band associates (not necessarily a visible band) to a separate primary color results in a color composite image. This research used false color composite with 652 RGB bands and is combined using ArcGIS 10.5 software.

- 3) Pan-sharpening
Pan-sharpening is a technique that merges high-resolution panchromatic data with medium-resolution multispectral data to create a multispectral image with high resolution features (Jaya, 2010). The Brovey transformation is based on spectral modeling and was developed to increase the visual contrast in the high and low ends of the data's histogram (ESRI, 2005). It uses a method that multiplies each resampled multispectral pixel by the ratio of the corresponding panchromatic pixel intensity to the sum of all the multispectral intensities. Pan-sharpening is processed using ArcGIS 10.5 software.
- 4) Classification of Land Use (Maximum Likelihood Method)
A maximum likelihood classification algorithm is one of the well-known parametric classifiers used for supervised classification. According to (ERDAS, 1999), the weighted distance or likelihood D of unknown measurement vector X

belong to one of the known classes M_c is computed based on the Bayesian equation:

$$D = \ln(a_c) - [0.5 \ln(\%Cov_c\%) - [0.5(X-M_c)^T (Cov_c^{-1} - 1) (X-M_c)]] \quad (1)$$

where:

D = weighted distance (likelihood)

c = a particular class

X = unclassified vector based

M_c = predetermined class

a_c = percent probability that any candidate pixel is a member of class

c (defaults to 1.0, or is entered from a priori knowledge)

Cov_c = the covariance matrix of the pixels in the sample of class c

$|Cov_c|$ = determinant of Cov_c (matrix algebra)

Cov_c^{-1} = inverse of Cov_c (matrix algebra)

T = transposition function (matrix algebra)

5) Ground Check

Ground check activities were conducted by using random sampling method proportionally of area of each class. Determination of coordinates of observation points in accordance with existing land use objects accompanied by observation of objects and imagery of land use visibility on the actual conditions in the ground check. In addition, topographic conditions and accessibility also support for point coordinate retrieval. Coordinate point taken in ground check as much as 86 points. This observation aims to match the land use that has been interpreted in the imagery visually with actual land use conditions in the field. Intake of coordinates of ground check data points using Geographic Positioning System (GPS).

6) Accuracy Assessment

Accuracy analysis was conducted to determine the accuracy of the classification made. Accuracy is analyzed using a contingency matrix that is a square matrix containing the number of classified pixels, shown in the table. This matrix is often called "error matrix" or "confusion matrix" (Stehman, 1997). In this contingency matrix, the analyst can also calculate the accuracy of the producer (accuracy) and the accuracy of the users (user's accuracy) of each class. The producer's accuracy is the accuracy obtained by dividing the correct pixels by the total number of pixels of sample regions per class. The equations used to measure the estimated overall accuracy, user's accuracy, producer's accuracy and kappa coefficient are as follows:

$$\text{Overall Accuracy} = \frac{\sum_{i=1}^r X_{ii}}{N} \times 100\% \quad (2)$$

$$\text{User's Accuracy} = (X_{ii} / X_{i+}) \times 100\% \quad (3)$$

$$\text{Producer's Accuracy} = (X_{ii} / X_{i+}) \times 100\% \quad (4)$$

$$\text{Kappa (K)} = \frac{n \sum_{i=1}^r X_{ii} - \sum_{i=1}^r (X_{i+} * X_{+i})}{n^2 - \sum_{i=1}^r (X_{i+} * X_{+i})} \times 100\% \quad (5)$$

where:

r = number of row and column in cross classification table,

n = number of pixels in the example,

X_{ii} = number of combinations along the diagonal,

X_{i+} = observations in row i , and

X_{+i} = observations in column i .

B. Evaluation of Land Use

Evaluation of land use suitability is used for current revision of space utilization allocation. Evaluation of land use suitability is conducted by comparing land use with regional spatial plan (RTRW). In this process, there is an overlay between the land use map the regional spatial plan (RTRW) of Geopark map. The result will be shown in map and table.

3. RESULT AND DISCUSSION

A. Land Use Map

Using GIS analysis, it is obtained the area and percentage of each classes land use in the research area such as Figure 3 and Table 1.




Figure 3.
Land Use Map of Batur UNESCO Global Geopark

Table 1 shown area of each classification in Batur UNESCO Global Geopark area is dominated by plantation with area of 7,317.60 ha (37.68%), followed by forest 3,901.22 ha (20.09%), mixed plantations 2,107.22 ha (10.85%), horticultural 1,736.43 ha (8.94%), Batur lake 1,532.09 ha (7.89%), bare land 1,003.63 ha (5.17%), bushes 933.12 ha (4.80%), built-up/residential 430.36 ha (2.22%), cloud 283.67 ha (1.46%), sand 112.84 ha (0.58%) and fish cultivation 64.21 ha (0.33%).

Table 1. Area and Percentage Based on Classification Results

No	Classes of land use	Area (ha)	Percentage
1	Built-up/Residential	430.36	2.22%
2	Forest	3,901.22	20.09%
3	Mixed plantations	2,107.22	10.85%
4	Horticultural	1,736.43	8.94%
5	Plantation	7,317.60	37.68%
6	Bushes	933.12	4.80%
7	Bare land	1,003.63	5.17%
8	Sand	112.84	0.58%
9	Batur lake	1,532.09	7.89%
10	Fish cultivation	64.21	0.33%
11	Cloud	283.67	1.46%
Total		19,422.39	100.00

Table 2. Confusion Matrix

GPS Data Clasifi- cation data	GPS Data										Row Total
	A	B	C	D	E	F	G	H	I	J	
A	3	0	0	0	0	0	0	0	0	0	3
B	0	15	0	1	0	0	0	0	0	0	16
C	0	1	6	1	2	0	0	0	0	0	10
D	1	0	0	5	1	0	0	0	0	0	7
E	0	0	3	0	25	0	0	0	0	0	28
F	0	0	0	0	1	4	0	0	0	0	5
G	0	0	0	0	0	0	4	0	0	0	4
H	0	0	0	0	0	0	0	3	0	0	3
I	0	0	0	0	0	0	0	0	6	0	6
J	0	0	0	0	0	0	0	0	0	3	3
Column Total	4	16	9	7	29	4	4	3	6	3	86

Overall accuracy: $(3+15+6+5+25+4+4+3+6+3)/86 = 86,05\%$

Where: A: Built-up/Residential
B: Forest,
C: Mixed plantations
D: Horticultural
E: Plantation
F: Bushes
G: Barren land
H: Sand
I: Batur lake
J: Fish cultivation

Table 3. Assessment Classification Results

No	Classification	Producer's Accuracy	User Accuracy
1	Built-up/Residential	3/4 = 75%	3/3 = 100%
2	Forest	15/16 = 93.75%	15/16 = 93.75%
3	Mixed plantations	6/9 = 66.67%	6/10 = 60%
4	Horticultural	5/7 = 71.43%	5/7 = 74.43%
5	Plantation	25/29 = 86.21%	25/28 = 89.29%
6	Bushes	4/4 = 100%	4/5 = 80%
7	Bare land	4/4 = 100%	4/4 = 100%
8	Sand	3/3 = 100%	3/3 = 100%
9	Water body	6/6 = 100%	6/6 = 100%
10	Fish cultivation	3/3 = 100%	3/3 = 100%

The method used to calculate the accuracy of classification by using the error matrix is then done producer's accuracy calculation, user accuracy,

overall accuracy and kappa coefficient value (Table 2 and Table 3).

The accuracy in Table 3 value for mixed plantation, horticultural, and plantation has a value below 100%, this is due to the small area of land ownership for each land use. Plantation and horticultural lands are being planted by other trees as barriers and shade trees to be detected as mixed plantation. From Table 3 if calculated value of Kappa accuracy based on the equation of number 5 obtained value 0.83.

Calculation of the overall accuracy shows that the process of map classification is acceptable because it has an overall accuracy value of more than 85% (Anderson, 1976). Meanwhile, according to Landis and Koch (1977), the coefficient value of kappa 0.81-1 is

included in very high category, this means that the value of the accuracy test on the classification results can be trusted.

B. Evaluation of Land Use with Regional Spatial Plan

Evaluation of land use with regional spatial plan is done to determine the suitability of land use utilization with the planning documents that has been established in regional regulation. With overlay method, the level of suitability for each land use in Batur UNESCO Global Geopark area will be known. Below are the results of the evaluation (Figure 10 and Table 4).

Overall Table 5 shows the land use in Batur UNESCO Global Geopark has deviations with regulation of regional spatial plan (RTRW). Overall




Figure 4.

Map of Evaluation Existing Land Use with Spatial Plan

percentage of suitable area is 10,863.14 ha (55.93%) and not suitable 8,275.58 ha (42.61%) with cloud interference/not detected 283.67 ha (1.46%). To find out more detailed comparison between spatial plans with land use for built-up/residential will be shown in Figure 5 and Figure 6.

Intersect overlay between the land use (Figure 7) with spatial plan resulted suitable only 114.64 ha or 26.64% of total built-up/residential. So the built-up/residential in Batur UNESCO Global Geopark 73.36% is not suitable with the regional spatial plan.

Protection of the forests need to be improved in this region, from Figure 12 shown forest condition in Batur UNESCO Global Geopark is very different from the spatial plan. The analysis results in Table 4 shows that only 2,587.98ha (66.34%) suitable with the spatial plan. Figure 8-11 are show the forest area in spatial plan.

Table 4. Result Evaluation Land Use with Regional Spatial Plan

No	Land Use	Spatial Plan											Total
		A	B	C	D	E	F	G	H	I	J	K	
1	Built-up/ Residential	50.68	63.96	13.87	19.73	49.72	52.24	167.88	0.04	7.42	-	4.82	430.36
2	Forest	5.17	15.62	1,751.40	61.32	273.04	502.22	1,266.82	6.17	10.63	-	8.83	3,901.22
3	Mixed Plantations	14.13	43.10	453.56	54.97	222.57	126.09	1,175.30	3.57	4.14	-	9.79	2,107.22
4	Horticultural	37.47	336.30	134.96	65.85	153.01	119.25	833.49	8.84	28.80	-	18.46	1,736.43
5	Plantation	55.60	119.75	1,865.92	160.20	784.93	356.81	3,911.95	33.61	7.43	-	21.40	7,317.60
6	Bushes	2.85	22.51	12.07	88.19	25.84	719.73	55.48	3.92	-	-	2.53	933.12
7	Bare land	1.05	0.36	118.21	4.73	24.23	716.09	84.54	1.60	17.47	-	35.35	1,003.63
8	Sand	1.87	3.49	0.46	1.84	64.00	15.22	21.79	-	1.70	-	2.47	112.84
9	Batur Lake	-	-	-	-	-	-	-	-	17.09	63.35	1,451.65	1,532.09
10	Fish Cultivation	-	-	-	-	-	-	-	-	2.74	1.92	59.55	64.21
11	Cloud	9.11	3.45	35.03	0.41	21.52	1.43	206.45	6.09	0.09	-	0.09	283.67
Total		177.93	608.54	4,385.48	457.24	1,618.86	2,609.08	7,723.70	59.92	101.43	65.27	1,614.94	19,422.39

A : Built-up/ residential	B : Special Tourism Object	C : Protected Forest
D : Limited Production Forest	E : Grand Forest Park	F : Nature Park Tourism
G : Plantation	H : Horticultural	I : Lake Border
J : Fish Cultivation	K : Lake	: Suitable
: Not Suitable	: Cloud/Not detect	

Table 5. Percentage of Suitable Area between Land Use with Regional Spatial Plan

No	Suitability	Area (ha)	Percentage (%)
1	Suitable	10,863.14	55.93
2	Not suitable	8,275.58	42.61
3	Cloud/Not detected	283.67	1.46
Total		19,422.39	100.00

Land use for plantations is scattered in the region of Batur UNESCO Global Geopark, considering Bangli Regency is a producer of Kintamani arabica coffee. This plantation is also interspersed with other tree like orange tree (*citrus sinensis*, *citrus nobilis*, *citrus reticulata*, *citrus unshiu*), mangoes tree (*mangifera L.*), cashew tree (*anacardium occidentale*), clove tree (*syzygium aromaticum*), jack tree (*artocarpus heterophyllus*). The analysis results obtained the land for plantation and mixed



Figure 5
Plan of Built-up/Residential



Figure 6
Plan of Special Tourism Object




Figure 7.
Land use for Built-up/Residential



Figure 8.
Plan of Protected Forest



Figure 9.
Plan of Limited Production Forest



Figure 10.
Plan of Grand Forest Park



Figure 11.
Plan of Nature Tourism Park




Figure 12.
Land Use for Forest

plantations in this region 7,317.60 ha and 2,107.22 ha. Land use for plantation in the spatial plan layout cover an area 7,723.70 ha, but if intersect overlay with land use is obtain the suitable only 5,087.25 ha (53.98%). Figures 14 and Figure 15 showed the spread of land use for plantation and mixed plantations increase of the plan (Figure 13).



Figure 13.
Plan of Plantation




Figure 14.
Land Use for Plantation



Figure 15.
Land Use for Mixed Plantation

Land use for horticultural also increase significantly. From the analysis, land use for horticultural resulted 1,736.43 ha from 59.92 ha in the spatial plan document. Land use for horticultural in spatial plan document located in the west of Kintamani district, but in fact the development of horticultural takes place in the east of Kintamani district (Batur UNESCO Global Geopark). The analysis result using intersect overlay land use with spatial plan obtains suitable 8.84 ha or only 0.51% of land use for horticultural, the remaining 99.49 % land use for horticultural not suitable with the spatial document. Figure 16 and 17 shows the plan and land use for horticultural.

Utilization of the lake for fish cultivation also increased. Fish cultivation in spatial plan located in north of the lake to northeast (Figure 18). In the spatial plan document fish cultivation allocation 65.27 ha, but the land use spread to the south and west of the lake with total area 64.21 ha (Figure 5.19). Overlay result fish cultivation with spatial plan obtained suitable area only 1.92 ha (2.99%), the remaining 97.01 % fish cultivation not suitable with the spatial document.

Batur UNESCO Global Geopark has natural mineral resources. Sand detected by satellite images has an area 112.84 ha is mostly located in the northern part of Batur volcano (Figure 20). But the existence of sand in grand forest park and plantation area are mining by community and sold illegally (Figure 21). In other place of sand mined site produces large and deep holes. As conservation areas (Geopark) this activity is very contrary to the concept of



Figure 16.
Plan of Horticultural




Figure 17.
Land Use for Horticultural



Figure18.
Plan of Fish Cultivation



Figure 19.
Fish Cultivation

protection, education and sustainable development. Geopark is basically aimed at raising awareness and understanding of issues in society, such as sustainable use of natural resources, reducing the impacts of climate change and reducing the risk of natural disasters.



Figure 20.
Sand in Batur UNESCO Global Geopark



Figure 21.
Mining Activity in Batur UNESCO Global Geopark

4. CONCLITION AND SUGGESTION

Land use in Batur UNESCO Global Geopark area until 27 September 2017 interpretation with Landsat 8 satellite imagery, utilization the area in Batur UNESCO Global Geopark area is dominated by plantation with area of 7,317.60 ha (37.68%), followed by forest 3,901.22 ha (20.09%), mixed plantations 2,107.22 ha (10.85%), horticultural 1,736.43 ha

(8.94%), Batur lake 1,532.09 ha (7.89%), bare land 1,003.63 ha (5.17%), bushes 933.12 ha (4.80%), built-up/residential 430.36 ha (2.22%), sand 112.84 ha (0.58%) and fish cultivation 64.21 ha (0.33%). Result of evaluation land use with regional spatial plan (RTRW) in Batur UNESCO Global Geopark overall has suitable area 10,863.14 ha (55.93%), not suitable area 8,275.58 ha (42.61%) and not detected/cloud interference 283.67 ha (1.46%) of 19,422.39 ha total area.

Suggestions for research development and all stakeholders in Batur UNESCO Global Geopark area are: (1) need for further research on the causal factor of land use change in the area of Batur UNESCO Global Geopark, (2) need community awareness in Batur UNESCO Global Geopark area because many threats and problems caused when land use is not suitable with regional spatial plan, (3) to Bangli government, before preparing a Detailed Spatial Plan of Batur needs to be considered changes to the current region spatial plan because the deviation is already very high with the existing land use, and (4) to central and local governments in preparing the master plan in the UNESCO Global Geopark Batur area should pay attention to land capacity and land functions.

REFERENCES

- Adnyana, I W.S., Nishio, F., Sumantyo, J.T.S., Hendrawan, G. 2006. Monitoring of Land Use Change Using Aerial Photograph and Ikonos Image in Bedugul, Bali. *Remote Sensing and Earth Science* 3: 51-57.
- Agustini, N.L.P.K. 2015. "Destination Management and Imaging Kintamani Towards Sustainable Tourism" (*Dissertation*) Denpasar: Udayana University. Available from: <http://erepo.unud.ac.id/10100/1/4f6bbe35ad8d7260e6cb25964432a49a.pdf>.
- Anderson, J.R., Hardy, E.E., Roach, J.T., Witmer, R.E., 1976 *Land Use and Land Cover Classification System for Use with Remote Sensor Data*. Washington: Geological Survey Professional Paper 964. Available from: <https://pubs.usgs.gov/pp/0964/report.pdf>.
- Bangli Government. 2013. *Peraturan Daerah Kabupaten Bangli Nomor 9 Tahun 2013 tentang Rencana Tata Ruang Wilayah Kabupaten Bangli 2013-2033*. Bangli: Government of Bangli Regency.
- ERDAS. 1999. *Field Guide Fifth Edition, Revised and Expanded*. US: ERDAS Worldwide Headquarters. Available from: <http://web.pdx.edu/~emch/ip1/FieldGuide.pdf>.
- ESRI. 2005. *GIS Solutions for Environmental Management*. New York: Environmental

- Systems Research Institute, Inc. Available from: <http://www.esri.com/library/brochures/pdfs/gis-sols-for-env-mgmt.pdf>
- Hidayati, I. N., Toyibullah Y. 2011. Kajian Indeks Potensi Lahan Terhadap Pemanfaatan Rencana Tata Ruang Wilayah Menggunakan Sistem Informasi Geografis di Kabupaten Sragen. *Globe*, 13 (No. 2): 156–164.
- Jaya, I N.S. 2010. *Analisis Citra Digital Perspektif Penginderaan Jauh untuk Pengelolaan Sumber Daya Alam*. Bogor: IPB Press.
- Knight, E.J., Kvaran, G. 2014. Landsat-8 Operational Land Imager Design, Characterization, and Performance. *Remote Sensing*, 6: 10286-10305.
- Landi, J., Koch, G. 1977. The Measurement of Observer Agreement for Categorical Data. *Biometrics* 33:159–74 available from https://www.dentalage.co.uk/wp-content/uploads/2014/09/landis_jr__koch_gg_1977_kappa_and_observer_agreement.pdf
- Naningsi, N. 2016. “Study of Land Use Change From 1997 to 2014 Using Landsat Data In Bangli Regency” (*tesis*). Denpasar: Udayana University.
- Richards, J.A. 2013. *Remote Sensing Digital Image Analysis: An Introduction*. 5th. New York: Springer Heidelberg.
- Stehman, S. 1997. Estimating Standard Errors of Accuracy Assessment Statistics Under Cluster Sampling. *Remote Sensing of Environment*, 60: 258–269.
- Wirosoedarmo, R., Widiatmono J.B.R., Widyoseno, Y. 2014. RTRW Arrangement Based on Environmental Supportability Based on Land Capability. *Agritech*, 34 (No. 4): 463-472. Available from: <https://media.neliti.com/media/publications/92932-ID-rencana-tata-ruang-wilayah-rtrw-berdasar.pdf>.