

## EFFECTIVENESS OF TEA TWIG LIQUID SMOKE (*Camellia sinensis*) AS AN ANTISEPTIC AGAINST MICROBES IN VIVO AND IN VITRO

Aulia Nur Arrsy<sup>1</sup>, Ayu Rahayu Nurul Ni'am<sup>1</sup>, Salsabila Zahroh<sup>1</sup>, Feldha Fadhila<sup>1</sup>, Nindya Sekar Mayuri<sup>2,3</sup>,  
Alfi Rumidatul<sup>2</sup>

<sup>1</sup> Department of Medical Laboratory Technology, Rajawali Institute of Health

<sup>2</sup>, School of Life Sciences and Technology ITB,

<sup>3</sup> Pharmacy Study Program, Politeknik Meta Industri

Correspondence email: [alfi@sith.itb.ac.id](mailto:alfi@sith.itb.ac.id)

### ABSTRACT

Transmission of disease-causing microorganisms to humans can occur directly or indirectly. Efforts to prevent disease transmission can use antiseptics, but the use of antiseptics has a number of side effects, therefore other alternatives are needed. The content of compounds in tea twig liquid smoke that can have potential as an antiseptic. The purpose of this research is as a means to see whether there is an effectiveness of tea twig liquid smoke on microbial growth in vitro which is used as an antiseptic with the help of an inhibition test and in vivo which is carried out by the swab test method on the palm of the hand. It can be seen from the results of the in vitro test to get the inhibition zone of tea twig liquid smoke (*Camellia sinensis*) against the growth of *Escherichia coli* ATCC 25922, *Staphylococcus aureus* ATCC 25923, *Aspergillus flavus* ATCC 9643, and *Candida albicans* 10231, the most effective results were obtained in grade 2 liquid smoke with a concentration of 75% with an inhibition zone diameter of 18.1 mm (*E. coli*), 15.6 mm (*E. coli*), 15.6 mm (*Candida albicans* 10231), 15.6 mm (*S. aureus*), 32.3 mm (*A. flavus*) and 4.3 mm (*C. albicans*). In vivo test results prove that there is an effectiveness of reducing the amount of growth in bacterial colonies on average by 88.33%, while the results of the effectiveness of reducing the number of fungal colonies on average by 91.52%. From the overall questionnaire data, it was found that 66% of respondents liked the aroma, 70% liked the color, 92% liked the dryness, and as many as 92% of respondents did not cause samaping effects on the skin such as itching, heat or burning from 75% tea twig liquid smoke. Through this study, the research results were obtained in the form of conclusions explaining that tea twig liquid smoke (*Camellia sinensis*) has the potential as an antiseptic.

**Keywords:** Liquid smoke., tea twigs., antiseptic

### INTRODUCTION

Transmission of disease-causing microorganisms to humans can occur directly or indirectly. Microorganisms that can cause infection are bacteria and fungi. *Escherichia coli* is known as one of the bacteria that cause intestinal infections, especially diarrhea.<sup>1</sup> According to the Indonesian Ministry of Health, Extraordinary Events (KLB) to be precise in 2017 there have been records of diarrheal diseases occurring 21 times which have spread to 12 provinces and 17 districts / cities with 1725 patients and 34 deaths.<sup>16</sup> Another bacterium that can cause infection is *Staphylococcus aureus* which is a trigger for nosocomial infections. Indonesia itself has a nosocomial infection prevalence record of 7.1%.<sup>3</sup> *Staphylococcus aureus* can also be the cause of cases of Acute Respiratory Infection (ARI). In a study by Rudan et al, (2021) stated that the latest cases of ARI in the world amounted to 156 million per year. Among these numbers, 151 million cases (96.7%) were found in developing countries. If ARI treatment is not

carried out properly, it can lead to complications that are classified as serious, such as the emergence of infections in the lungs, infections in the membranes of the brain, decreased awareness, respiratory failure, and even death, especially in infants and toddlers who do not have optimal immune strength.<sup>4</sup> Aspergillosis is the term refers to diseases of the respiratory system caused by the presence of fungal infections of the genus *Aspergillus*. Specialist causes are cosmopolitan including *Aspergillus flavus* (*A. flavus*).<sup>11</sup> The Medical Records Installation of Dr. Sarjito Hospital said there were 902 cases of otomycosis found in the city of Yogyakarta between 2014 and December 2018. Otomycosis disease is caused by the growth of foreign substances from the genus *Aspergillus*. One of the fungal infections that has been widely spread in Indonesia is Candidiasis. The main cause of the fungal infection is the agent *Candida* sp.<sup>31</sup> A number of epidemiological studies conducted in Hong Kong explained that the number of candidiasis incidence rates found in the Asian region was found to originate from *C. albicans* which is a species

most often known to be the cause with an average of 56% of candidiasis cases. *C. albicans* as the dominant species has had incidence rates ranging from 37% in Latin America to 70% in Norway. This has been the result of an increase in the incidence of invasive candidiasis along with an increase in the population of susceptible individuals, and the hindrance of treatment by antifungal resistance.<sup>19</sup>

Efforts to prevent disease transmission can be done with the habit of washing hands using soap.<sup>12</sup> The results of global research show that the use of soap for hand washing habits can not only minimize the level of disease transmission but can also be useful as a prevention of the incidence of diarrheal diseases by 50% and ARI by 45% (Nurdin, 2018), according to Yannuarist et al. (2020) alcohol, glycerol, tricholosan and other chemicals contained in hand sanitizers can cause side effects in the form of dryness and irritation located on the skin which generally occurs in children. Therefore, an alternative to natural ingredients is needed as a substitute for hand sanitizer ingredients, one of which is tea twigs which has the potential as an anti-microbial. Green tea stores the largest content in the form of flavonoid compounds which are referred to as a class of polyphenolic compounds.<sup>22</sup> Referring to the results of phytochemical tests conducted using ethanol solvents, it is stated that tea twigs contain flavonoid compounds and catechins.<sup>13</sup> It can be known that flavonoids are active compounds that can be utilized as antibacterial, anti-inflammatory, and even antifungal. The process of flavonoids in inhibiting fungal growth can trigger disruption of fungal cell membrane permeability.<sup>5</sup>

The content of compounds in tea twigs can be obtained by pyrolysis method, the result of the pyrolysis process used is liquid smoke. The term liquid smoke can be called liquid smoke which is intended as a result of condensation or condensation from the evaporation process from combustion carried out either directly or indirectly derived from various materials that contain lignin, cellulose, hemicellulose and other carbon compounds that can bring various compounds such as phenols, carbonyls and so on. From several studies, it is said that liquid smoke is used as anti-bacterial, anti-fungal, anti-insect, and anti-termite.<sup>24</sup> Liquid smoke is unique in that it can have antibacterial properties. The existence of properties that function as antibacterial has a relationship with the content of phenol and acetic acid contained in liquid smoke. The ability of phenol compounds is believed to cause damage to the cytoplasmic membrane resulting in leakage of the membrane which can interfere with bacterial growth and can even result in death. In addition, acetic acid can also cause destabilization of various functions and structures in cell components.<sup>30</sup> Meanwhile, the results of research by Lestari (2017), show that the extract produced from tea leaves can be used as an inhibitor of *Aspergillus flavus* growth. Based on the description above, it shows that tea twig liquid smoke has the potential to inhibit the growth of bacteria and fungi, so further research needs to be done so

that the benefits of tea twig liquid smoke inhibit the growth of bacteria and fungi can be known in more detail. The purpose of this study was conducted as a means to see whether there is an effectiveness of tea twig liquid smoke on microbial growth in vitro which is used as an antiseptic with the help of an inhibition test and in vivo which is carried out by the swab test method on the palm of the hand.

## MATERIALS AND METHODS

The method used in this study is to use an experimental descriptive method, and carry out testing at the Microbiology Laboratory of the Rajawali Institute of Health Bandung from January to February 2022 by means of in vitro and in vivo. In the implementation of this study, the samples used in the in vitro test were *Escherichia coli* ATCC 25922, *Candida albicans* ATCC 10231, *Staphylococcus aureus* ATCC 25923, and *Aspergillus flavus* ATCC 9643 tested with tea twig liquid smoke (*Camellia sinensis*) at concentrations of 35%, 50%, and 75%. The number of samples used in in vivo testing amounted to 24 respondents consisting of men and women, each of which amounted to 12 respondents. Related to the Swab test through the palm of the respondent is carried out before the start and after the action is given to the liquid smoke of the optimum concentration of tea twigs from the in vitro test. In determining the number of respondents determined using the federer formula.

### Research tools and material

The research used various tools, namely erlenmeyer, autoclave, oven, petri dish, test tube, stirring rod, cotton swab, paper disc, micropipette, vernier, colony counter, and pH meter. The materials used were grade 1 and grade 2 tea twig liquid smoke, 70% alcohol, sterile distilled water, nutrient agar medium, potato dextrose agar medium, 1% orange scent perfume, gram staining reagent, lactophenol cottonblue staining reagent, *Candida albicans* ATCC 10231, *Escherichia coli* ATCC 25922, *Staphylococcus aureus* ATCC 25923, and *Aspergillus flavus* ATCC 9643 cultures.

### Data processing and analysis


Effectiveness results from a decrease in the number of microbes. Questionnaire data in the form of secondary data obtained from respondents with several questionnaire parameters including preference for color, aroma, dryness effect, itching effect and burning. The questionnaire assessment used has several points, namely 0 for dislike, 1 is normal, 2 likes, and 3 is very like. There are criteria for respondents intended in this study, namely focusing on respondents who have not washed their hands using water or soap, do not have allergies to hand sanitizer or alcohol, do not have wounds on their hands, hand parts, and have the willingness to fill out a questionnaire sheet. The results obtained are in the form of calculating the number of microbial colonies that grow on NA and PDA medium.

Measurements were made by comparing the growth of the number of microbial colonies from swab samples before and after being given 75% tea twig liquid smoke natural ingredient hand sanitizer. Then compared with the control results, namely using an antiseptic containing 70% alcohol. Then the data obtained is processed using the help of a computer program, namely Microsoft excel to determine the correlation results of the use of 75% tea twig liquid smoke hand sanitizer.

## RESULTS

The observation of the characteristics of the tea twig liquid smoke solution is macroscopically seen in terms of color, which is yellowish brown with a pungent smoke aroma, and the consistency of the solution is liquid and has a pH of 2.6.

**Table 1.** Macroscopic Characteristics of Tea Twig (*Camellia sinensis*) Liquid Smo

Physical Properties	Liquid smoke from tea twigs 75% concentration	Picture
Color	Yellowish brown	
Aroma	Stinging	
Consistency	Liquid	
Clarity	Somewhat clear	
pH	2,6	


## Microbial Identification

**Table 2:** Microbial Identification

No	Type of Microbe	Colony Morphology	Cell Morphology
1	<i>Escherichia coli</i> ATCC 25922	Colonies are small round in shape, convex in elevation, smooth in texture and milky white/opaque in color	Short, red rods are Gram negative
2	<i>Staphylococcus aureus</i> ATCC 25923	The colonies are round and resemble grapes in shape, convex in elevation, and white in color	Round and clustered like grapes and purple are Gram positive
3	<i>Aspergillus flavus</i> ATCC 9643	Colonies are green to yellowish green with a granular and compact colony shape	Round to oval vesicles, round conidia, conidiophores long, cylindrical
4	<i>Candida albicans</i> ATCC 10231	Colonies are small round in shape white mucoid texture with smooth and even edges	Oval in shape, purple in color, Gram positive

Microbial identification was carried out on *Escherichia coli* ATCC 25922, *Candida albicans* ATCC 10231, *Staphylococcus aureus* ATCC 25923, and *Aspergillus flavus* ATCC 9643 carried out macroscopically and microscopically the results can be seen in table 2.

## In Vitro Test Results



**Figure 1.** Zone of Inhibition of Liquid Smoke Tea Twigs (*Camellia sinensis*) Grade 1 on Tet Microbes

Based on Figure 1, the results of the in vitro test showed that grade 1 liquid smoke of tea twigs (*Camellia sinensis*) can inhibit the growth of *E. coli* ATCC 25922 with a concentration of 35% with a size of 10.3 mm, 50% concentration with a size of 13.8 mm, and 75% concentration with a size of 17.6 mm. In vitro test results on *S. aureus* ATCC 25923, 35% concentration with a size of 5.5 mm, 50% concentration with a size of 9.1 mm, and 75% concentration with a size of 15.8 mm using the disc diffusion method. Then the results of

inhibition on the growth of *A. flavus* ATCC 9643 35% concentration with a size of 21 mm, 50% concentration with a size of 25.6 mm, and 75% concentration with a size of 27.5 mm. In vitro test results on *C. albicans* ATCC 10231 35% concentration with a size of 1.1 mm, 50% concentration with a size of 1.3 mm, and 75% concentration with a size of 5.6 mm using the well diffusion method.




Figure 2. Zone of Inhibition of Liquid Smoke Tea Twigs (*Camellia sinensis*) Grade 2 on Test Microbes

Figure 2 shows the results of the in vitro test of liquid smoke of tea twigs (*Camellia sinensis*) grade 2 can inhibit the growth of *E. coli* ATCC 25922 35% concentration with a size of 8.5 mm, 50% concentration with a size of 13.3 mm, and 75% concentration with a size of 18.1 mm. In vitro test results on *S. aureus* ATCC 25923 35% concentration with a size of 7 mm, 50% concentration with a size of 10.6 mm, and 75% concentration with a size of 15.6 mm using the disc diffusion method. Then the results

of inhibition on the growth of *A. flavus* ATCC 9643 35% concentration with a size of 26 mm, 50% concentration with a size of 27 mm, and 75% concentration with a size of 32.3 mm. In vitro test results on *C. albicans* ATCC 10231 35% concentration with a size of 2.3 mm, 50% concentration with a size of 3.5 mm, and 75% concentration with a size of 4.3 mm using the well diffusion method.

### In Vivo Test Results




Figure 3. Liquid Smoke Effectiveness TeaTwigs

Based on Figure 3, it shows the results of the effectiveness of 75% tea twig liquid smoke in inhibiting microbial growth with the in vivo method, and the results obtained have an average percentage decrease in the


<http://ojs.unud.ac.id/index.php/eum>  
doi:10.24843.MU.2024.V13.i12.P06

microbial growth with the in vivo method, and the results obtained have an average percentage decrease in the

number of bacteria by 88.33% and fungi by 91, 25, this indicates that 75% concentration of teatwig liquid smoke is

able to inhibit microbial growth.

### Questionnaire Results



**Figure 4.** Percentage of Smoke Characteristics Characteristics of Liquid Tea Twigs Overall Preferred Respondents

Questionnaire data collection is carried out with the aim of measuring the level of general public preference as a correspondent to tea twig liquid smoke products that are used as ingredients for making antiseptics. Based on Figure 4, the results show that 66% do not mind the aroma of tea twig liquid smoke which has been added with 1% citrus aroma perfume, the addition of this perfume aims to disguise the aroma like burnt results but 34% of respondents do not like the aroma of antiseptic made from 75% tea twig liquid smoke, this is because it can still smell smoke after a while after use and mixed with citrus fragrance which begins to disappear.

### DISCUSSION

Based on the results of measuring the inhibition zone in the liquid smoke of the twig (*Camellia sinensis*) with concentrations of 35%, 50%, and 75% can inhibit the growth of *A. flavus* ATCC 9643 and *C. albicans* ATCC 10231. The inhibition zone contained in each concentration is due to the presence of phenol compounds that can cause antifungal effects in tea twig liquid smoke because if the higher levels of phenol in liquid smoke cause its antifungal activity to increase as well. In inhibiting fungal growth, phenol denatures cell proteins and damages cell membranes, so do not be surprised if it causes death to pathogens.<sup>20</sup> Flavonoid compounds have a role as antifungals aimed at forming a complex compound and forming extracellular proteins and soluble proteins that cause disruption of fungal cell membrane permeability. Not only that, the ability of flavonoids can also inhibit the work of enzymes that have a role in the synthesis of fungal cell walls such as mannansynthase, chitin synthase, and glucan synthase. If there are inhibitors when these enzymes work, it will certainly interfere with the process of forming fungal cell walls and inhibit the process of

pseudohyphae formation.<sup>21</sup> As for the results of the measurement of the inhibition zone against liquid smoke of tea twigs (*Camellia sinensis*) with concentrations of 35%, 50%, and 75% can inhibit the growth of *E. coli* ATCC 25922 and *S. aureus* ATCC 25923. The occurrence of the inhibition zone is caused by the presence of phenol compounds that can inhibit bacterial growth in low concentrations of phenol can cause damage to the cytoplasmic membrane. And this damage can be an opportunity for organic ions, nucleotides and amino acids to also leave the cell and will ward off the entry of various necessary materials into the cytoplasmic membrane cells do not function properly.<sup>15</sup>

The results of the analysis of the characteristics of tea twig liquid smoke obtained that grade 2 tea twig liquid smoke has a slightly cloudy clear yellow color and has better antifungal properties than grade 1 tea twig liquid smoke because it contains very high acids and phenols and has a very low pH level value compared to grade 1 tea twig liquid smoke. Grade 1 tea twig liquid smoke has a clear yellow color due to the separation of harmful compounds and high tar in tea twig liquid smoke (*Camellia sinensis*). It is explained that the pH of good liquid smoke is in the range of 1.5-3.7 because in low pH conditions fungi with spores cannot live and multiply so that they can potentially inhibit the growth rate of fungi.<sup>26</sup> The content of active compounds contained in liquid smoke consists of phenol compounds and organic acids with anti-microbial abilities.<sup>9</sup> In addition to phenol, liquid smoke also contains other compounds in the form of acetic acid. The acidic antimicrobial effect of liquid smoke may directly shape a cytoplasm to be acidic, damaging the surface tension of the membrane and the loss of active transport of food through the membrane, causing destabilization of many types of functions and structures of cell components.<sup>20</sup> Acidic



compounds can also interfere with bacterial cell metabolism by lowering the pH of bacterial cells.

Phenol compounds contained in liquid smoke have the ability to damage the protein structure of bacteria by means of hydrogen bonds, so that bacterial metabolic activity will stop.<sup>10</sup> According to Aisyah (2019), the mechanism of alcohol compounds in inhibiting microbial growth, namely that in bacteria alcohol compounds react with lipid proteins, then will denature the proteins that make up the cell membrane and bacterial cell wall, while in fungi it will denature the fungal conidia membrane and dissolve the lipids that make up the hyphal cytoplasmic membrane and hyphal cell wall. Acidic compounds contained in liquid smoke can break through the cell walls of microorganisms, causing the cells of microorganisms to lysis. Tea twig liquid smoke with a concentration of 75% has the ability to inhibit microbial growth. Based on Kondo's research (2017), it states that the higher the concentration level in the liquid smoke solution, the stronger the ability to inhibit bacterial growth. Supported by Mayasari's research (2022), liquid smoke without purification or without dilution has strong inhibition against three types of bacteria, namely *Escherichia coli*, *Salmonella sp.*, and *Staphylococcus aureus*. In a study conducted by Aznury (2021), that people in choosing hand sanitizers prefer hand sanitizers with a very non-pungent aroma, this is because people do not really like the smell of hand sanitizers that are very pungent.

Most respondents were interested in the color of the 75% concentration of tea twig liquid smoke, because it was yellowish brown, slightly transparent. This is supported by the research of Aznury et al. (2020), that the clear transparent color of the hand sanitizer gel is the most favorite color choice, this is because it has attractiveness and is brighter in terms of color, while the dark color is less attractive to the public. Color selection for a product has an important role in order to be of interest to many people.<sup>29</sup> It is known that most people feel like the dryness caused after the use of 75% tea twig liquid smoke and only a small percentage feel dislike the dryness of 75% tea twig liquid smoke. People feel more comfortable when using products that provide a cool feeling and are not too sticky when not in use, and prefer hand sanitizers with gel texture compared to liquid texture.<sup>6</sup> Figure 4 shows that most people 92% like the sensation effect caused, because they do not get side effects from using 75% tea twig liquid smoke hand sanitizer. Based on the color and aroma arising from the liquid smoke of tea twigs, it is still not possible as an antiseptic that people like. Aznury et al. (2020) stated, if the use of hand sanitizer products is sticky, feels hot, causes irritation, it will interfere with the health of its users. So that the comfort factor of using hand sanitizer products must be prioritized such as in terms of color, aroma, texture, and the absence of side effects caused on the skin.

## CONCLUSIONS AND SUGGESTIONS

Based on the results of research that has been done, it can be concluded that grade 2 tea twig liquid smoke with a concentration of 75% has a better ability to inhibit the growth of bacteria and fungi.

## ACKNOWLEDGMENTS

For the implementation of this research, the author expresses his gratitude to various parties, especially those who contributed to the publication of this journal.

## REFERENCE

1. Agustine, L., Okfrianti Y., Jumiayati. (2018). Total Lactic Acid Bacteria Lactic Acid Bacteria (LAB) in Yoghurt with Sucrose and Skim Milk Variations. *Journal of World Nutrition*, 1(2), 79-83.
2. Aisyah, I., Giyanto, Sinaga, M. S., Nawangsih, A. A., Pari, G. (2018). *In Vitro* Test of Liquid Smoke against *Ralstonia syzygii subsp. Celebesensis*, the cause of Blood Borne Disease of Banana. *Journal Phytopatol Indonesia*, 14(4), 145-151.
3. Amelia, R., Burhanuddin, N. (2018). Identification of bacteria *Staphylococcus aureus* with nosocomial infection on bed linen in the hospital room. Nosocomial infections on bed linen in the room post-surgery care room at Makassar City Hospital Labuang baju Makassar city. In Proceedings of National Seminar Synergy of Multidisciplinary Science Science and Technology, (Vol. 1, pp. 272-278).
4. Amila, Pardedei, J.A., Simanjuntak, G. V., Nadeak, Y. (2021). Improving Knowledge parents about the dangers of smoking in the house and prevention of ARI in toddlers. *Jukeshum. Journal of Community Service*, 1(2), 65-70.
5. Asngad, A., & Nopitasari, N. (2018). Quality of Hand Sanitizer Gel (Handsanitizer) from Banana Stem Extract Banana Stem Extract with the Addition of Alcohol, Triclosan And Different Glycerin Dosage. *Bioexperiment. Journal Biology Research*, 4(2):61-70.
6. Aznury, Martha, Sari, R. P. (2020). Gel Hand Sanitizer Products Made from Based on Green Betel Leaf Liquid Extract (*Piper Betle* Linn.) as an Antiseptic. *Kinetics*, 11(1),

- 27-35.
7. Aznury, M., & Serlina, A. (2021). Optimization of soap making formula with addition of green betel leaf extract (piper betle l). *Kinetics*, 12(1), 51-59.
  8. Desvita, H., Faisal, M., Suhendrayatna (2020). Preservation of meatballs with edible coating of chitosan dissolved in rice hull- based liquid smoke. *Heliyon*, 6(10), 1-10.
  9. Erlytasari, D. N., Wibisono, G., Hapsari, R. (2019). Effectiveness of Liquid Smoke of Various Concentrations as Disinfecting Dental Clinic Tools. *Diponegoro Medical Journal. Journal Diponegoro Medicine*, 8(4), 1114- 1123.
  10. Fauziati, F., & Sampepana, E. (2021). Study of Chemical Characteristics Liquid Smoke of Palm Kernel Shell, Empty Empty Palm Fruit Bunches as Antibacterial and its Application. *Journal Industrial Technology Research*, 15(2), 338-347.
  12. Hasanah, U. (2017). Get to know Aspergillois, a fungal infection of the genus *Aspergillus*. *Journal of Healthy Family Sejahtera*, 15(2):76-86.
  14. Huliatusunisa, Y., Alfath, M. D., Hendiati. D. (2020). Handwashing Using Soap. *Journal Post Dharma Community Service Community. Journal of Post Dharma Community Service* (2), 40-46.
  15. Hung, Y. C., Hsiao, Y. H., Hsieh, J. F. (2021). Catechin content and free radical scavenging activity of *Camellia sinensis* twig extracts. *International Food Research Journal*, 8(2).
  16. Hutasoit, Dion, P. (2020). Effect of food sanitation and *Escherichia coli* bacteria on Diarrhea disease. *Scientific Journal Sandi Husada Health*, 9(2), 779- 786.
  17. Iryani, Suryeita, Putri, D. (2013). Inhibition test of coir liquid smoke areca nut liquid smoke on the growth of bacteria *bacillus cereus* and *Pseudomonas aeruginosa*. *Periodic*, 2(1), 38-42.
  18. Indonesian Ministry of Health. (2019). Profile Indonesia Health Profile. Jakarta.
  19. Kondo, S. A., Wibisono, G., & Ciptaningtyas, V. R. (2017). Effect of liquid smoke application at various concentrations on growth of *streptococcus sanguis* which causes gingivitis. *Journal Diponegoro Medicine (Diponegoro Medical Journal)*, 6(1), 106-113.
  20. Lestari, P.I. (2017). Activity Antifungal Activity of Tea Leaf Extract Against the Growth of *Aspergillus flavus*. *The Indonesian Journal of Infectious Diseases*, 1(01), 29-38.
  21. Lim, C. S. Y., Rosli, R., Seow, H.F., Chong, PP. (2012). *Candida* and Invasive Candidiasis: Back to Basics. *Europe Journal of Clinical Microbiology & Infectious Diseases*, 31(1), 21-31.
  22. Mahmud, M. (2022). Quality of Bali Beef Muscle *Longissimus Dorsi* (Ld) Muscle That Is Marinated in Liquid Smoke Flour And *Moringa* Flour (Doctoral dissertation, Hasanuddin University). 5(3), 46-51.
  23. Mayasari, U. (2022). Activity of antibacterial activity of Pirdot Leaf extract (*Saurauia Vulcani* Korth) in inhibiting the growth of bacteria *Pseudomonas Aeruginosa* bacteria causing nosocomial pneumonia infection. *Journal of researchgate* vol. 4 No. 1, 2020, 1-5.
  24. Mukty, I., Wirjatmadi, B., Purwanto, B. (2018). Effect of Giving Green Tea (*Camellia sinensis*) on the Activity of Superoxide Dismutase (SOD) Effect of Green Tea (*Camellia sinensis*) Against Superoxyde Dismutase (SOD). *Scientific Journal of Medicine Wijaya Kusuma*, 7(2), 158-164.
  25. Nurdin, I., & Hidayat, F. R. (2018). Relationship between Nail Clipping and Handwashing with Soap Handwashing with Soap (HWWS) at School with the Incidence of Diarrhea in Class IX at SMPN 36 Kelurahan Rapak Dalam Samarinda Seberang 2018. 9-12.
  26. Oramahi, E. R. P., Wardoyo, Kustiati. (2018). Chemical Characterazation and Biological Activities of Wood Vinegar from Wood Wastes of Furniture Industry. *Journal of Wood Science*. 64(3), 311-317.

27. Praja, R.N. (2017). Isolation and Identification of *Aspergillus* spp. on Kampung Chicken Lungs that Sold in Banyuwangi Market. *Journal Veterinary Medicine*. Vol 1, 6-11.
28. Puji, L.T. (2015). Analysis of Chemical Physicochemical and Anti bacterial Properties of Smoke Oil Palm Shell Liquid Smoke for Food Preservative. *Journal of Research Industrial Technology*, 4(8), 1-8.
29. Rachmawati, D. P., Rabbani, K., Rumidatul, A., Fadhila, F., Maryana Y. (2020). Activity Testing Antimicrobial Activity of Bark and Wood Extracts Sengon (*Falcataria Moluccana*) Twigs With N-Solvent Hexane, Ethyl Acetate and Methanol Against Enterobacteriaceae, *Staphylococcus Aureus* And *Candida Albicans*. *Journal of Analytical Media Health*, 11(2), 70-82.
30. Saepul, A., Pitrianingsih S., Sodikin A. (2022). Effectiveness of Liquid Smoke from (*Coffe* sp.) as antiseptic against microbes in vitro and in vivo. *Medika Kartika. Journal of Medicine and Health*, 5(1), 21-33.
31. Setyawardhani, D. A., & Saputri, C. M. (2021). Manufacture and Test Organoleptic Test of Hand Sanitizer from Mango Leaf (*Mangifera indica*) with Maceration Method. *Equilibrium Journal of Chemical Engineering*, 4(1), 1-7.
32. Sumpono, S. (2018). Activity Test Antioxidant and Antibacterial Activity of Smoke Palm Kernel Shell Liquid. In *Proceedings of SNPS (National Seminar Science Education)*.171-178.
33. Widaty, S. Budimulja, U. (2016). Superficial Dermatormycosis. *Science Skin and Gender Diseases*. 7th Edition. 7. Publishing Board of FKUI. Jakarta.
34. Yannuarista, D., Rintania, S., Hilmi, M. (2020). Organoleptic Test and Effectiveness of Lime Extract as a natural hand sanitizer. In *Proceedings of the National Seminar on Applied Innovative Research (SENTRINOV)*, 6(1), 1127-1134.

