

## RELATIONSHIP BETWEEN PULMONARY VASCULAR THICKENINGS WITH CT-SEVERITY SCORE IN COVID-19-CONFIRMED PATIENTS

Alex Danu<sup>1</sup>, Sri Asriyani<sup>1</sup>, Nikmatia Latief<sup>1</sup>, Andi Alfian<sup>2</sup>, Harun Iskandar<sup>3</sup>, Rafikah Rauf<sup>1</sup>

<sup>1</sup>Radiology Department, Medical Faculty of Hasanuddin University, Makassar, Indonesia <sup>2</sup>Public Health Department, Medical Faculty of Hasanuddin University, Makassar, Indonesia <sup>3</sup>Pulmonology Department, Medical Faculty of Hasanuddin University, Makassar, Indonesia

**Corresponding author:** Sri Asriyani, Radiology Department, Medical Faculty of Hasanuddin University, Perintis Kemerdekaan KM 11 St., Tamalanrea, Makassar City, South Sulawesi, Indonesia, 90245. Phone : 0411-899-5513. [sri\\_asriyani@med.unhas.ac.id](mailto:sri_asriyani@med.unhas.ac.id)

### ABSTRACT

**Introduction:** Previous studies have shown pulmonary vascular thickening found in COVID-19 patients. In addition, the CT-Severity Score (CT-SS) scoring system have become a reference in assessing the severity of COVID-19. This study analyzes the relationship between pulmonary vascular thickening on CT-Scan thorax with CT-SS in patients with confirmed COVID-19. **Materials and Methods:** A retrospective cross-sectional study was conducted on 137 patients with confirmed COVID-19 at the Radiology Installation of Doctor Wahidin Sudirohusodo Hospital Makassar. Patients who were eligible and had undergone a chest CT-Scan were then measured for their pulmonary vascular thickness in the intralesional and extralesional areas of the secondary lung lobes. The CT-SS score was assessed to determine the degree of mild or severe disease based on the opacity of the involved lung segment. Independent T-test and Chi-square were used to determine the differences and relationships between the variables studied. **Results:** Based on CT-SS, most subjects were classified as mildly ill (CT-SS < 19.5, 78/137 people). The mean intralesional and extralesional vascular thickening was  $2.65 \pm 0.48$  and  $2.17 \pm 0.36$ , respectively. There was a significant difference between vascular thickness between subjects with mild and severe CT-SS categories (intralesion:  $2.4 \pm 0.33$  vs.  $2.99 \pm 0.47$ ,  $p=0.001$ ; extralesion:  $2.04 \pm 0.28$  vs.  $2.35 \pm 0.39$ ),

**Conclusion:** Vascular thickening on CT-Scan chest images is related to the severity of COVID-19 as measured by CT-SS scoring.

**Keywords:** COVID-19, Chest CT-Scan, *CT-Severity Score*, Pulmonary Vascular Thickening

### INTRODUCTION

COVID-19 (Coronavirus Disease 2019) is an infectious disease caused by a species of coronavirus, namely Severe Acute Respiratory Syndrome Coronavirus 2. SARS-CoV-2 has become a worldwide pandemic since it was first discovered in Wuhan, China, at the end of December 2019. The incidence rate until February 10, 2022, global COVID-19 cases has reached 404 million cases and caused 5,783 million deaths.<sup>1</sup>

One of the challenges regarding the disease that causes this pandemic is the difficulty of predicting the complications of respiratory failure caused. Abnormalities found on chest CT scan images effectively predict complications of respiratory failure and mortality in COVID-19 patients, including those who are seriously ill and require intensive care.<sup>2,3</sup> Pulmonary vascular thickening on CT-Scan Thorax is the main feature in patients with confirmed COVID-19 concurrently with pneumonia This vascular thickening is also found in 72-78% of patients with COVID-19 This vascular thickening is believed to be a unique feature of

COVID-19 pneumonia that has not been previously described and is thought to contribute to respiratory failure in COVID-19 pneumonia.<sup>2,5</sup> The COVID-19 severity scoring system based on the chest CT-Severity Score (CT-SS) has recently been introduced and studied extensively. Yang et al. found an association between higher CT-SS scores and more severe COVID-19 infection. Abbasi et al. found a higher mortality rate in patients with higher CT-SS.<sup>6,7</sup>

### Materials and Methods

#### Population and Sample

This study used a retrospective cross-sectional design and was approved by the Health Research Ethics Committee, Faculty of Medicine, Hasanuddin University . We took secondary data in medical records from 137 patients (74 males, 63 females, mean age  $49.72 \pm 14.31$  years) who had confirmed COVID-19 through SARS-Cov-2 RT-PCR examination and had undergone CT examination. - Thoracic scan measurement at Wahidin Sudirohusodo Hospital Makassar between July to September 2021. Patients with a history of previous

chronic diseases (e.g., pulmonary hypertension, pulmonary embolism, post-chemotherapy, COPD, malignancy, chronic liver disease, and post-surgery) have an infiltration pattern of lesions only in the form of consolidation in all segments or lobes of the lung, and having a pleural effusion that covers part of the lung lobes were excluded from this study.

#### Imaging Technique

CT Scan equipment brand Light speed VCT 64 Slice GE with the technique of taking Scan from the tip of the lung to the costophrenic angle with the following scan parameters: tube voltage 120 kV, tube current-exposure time 50-150 mAs, scan thickness 5 mm with reconstruction thickness 1.5mm.

#### Imaging Interpretation

Analysis of the characteristics of the patient's chest CT scan was performed on the lung window (width 1500 HU; level -700 HU) and mediastinal (width 350 HU; level 40 HU) to measure the vascular thickness. The mean values for intralesional and extralesional secondary pulmonary lobe vascular thickening were then calculated. Further confirmation will be carried out by 2 (two) radiology specialists who have more than 10 years of clinical experience.

#### CT Severity Score

The CT severity score (CT-SS) for assessing the severity of COVID-19 was adapted from the CT-SS by Yang et al. This assessment uses lung opacification in each segment to assess the extent of the lesion and is evaluated by giving a score of 0, 1, and 2 if parenchymal opacification involvement is found to be 0%, <50% or

50%. CT-SS is determined from the number of scoring for each segment, with a range of 0-40 points. Patients are categorized as mild if the total score is < 19.5 and severe if the total score is 19.5.

#### Statistical Analysis

Vascular thickness data are numerical measurement scales expressed in mean  $\pm$  standard deviation or median  $\pm$  interquartile range (IQR) if not normally distributed. Other categorical data, both demographic status, CT-SS expressed in percentage and frequency (n, percent). The Statistical Program for Social Science (SPSS) software version 23.0 (IBM, USA) was used to perform data analysis. An Independent T-test was used to find the mean intralesional and extralesional vascular thickness difference between the two categories obtained from CT-SS scoring. Then a Chi-square test was performed to analyze the relationship between CT-SS and. A significant value is stated if the p-value < 0.05 is obtained.

## RESULTS

#### Baseline Characteristics

The baseline characteristics of this study can be seen in Table 1. In the 137 subjects studied, the mean age of the patients was  $49.72 \pm 14.33$  years, with the majority aged 56-65 years (29.2%) followed by 46- 55 years (24.1%). In addition, the majority of patients were male (54%), had a CT-SS score < 19.5 (56.9%). The mean intralesional and extralesional vascular thickening measured on CT-Scan Thoracic results was  $2.65 \pm 0.48$  and  $2.17 \pm 0.36$ , respectively.

**Table 1:** Sample distribution based on baseline characteristics

Characteristic	n = 137	%
<b>Age (years)</b>	$49.72 \pm 14.331$	
0 - 5	0	0.0%
5 - 11	0	0.0%
12 - 16	2	1.5%
17 - 25	4	2.9%
26 - 35	21	15.3%
36 - 45	20	14.6%
46 - 55	33	24.1%
56 - 65	40	29.2%
> 65	17	12.4%
<b>Gender</b>		
Male	74	54
Female	63	46
<b>CT Severity Score</b>		
<19.5	78	56.9
$\geq$ 19.5	57	41.6
<b>Vascular thickening (mm)</b>		
Intralesion	$2.65 \pm 0.48$	
Extralesion	$2.17 \pm 0.36$	

**Table 2:** Comparison of intralesional and extralesional vascular thickness with CT Severity Score

Location	Mean ± Standard Deviation of Vascular Thickness		p
	CT-SS <19.5	CT-SS ≥19.5	
Intralesion	2.4 ± 0.33	2.99 ± 0.47	0.001
Extralesion	2.04 ± 0.28	2.35 ± 0.39	0.001

Independent T -test

**Intralesional and Extralesional Vascular Size Comparison Based on CT-SS Score**

Table 2 shows the results of comparing intralesional and extralesional vascular sizes based on CT-SS scoring. It was found that the mean intralesional vascular size in patients with CT-SS < 19.5 was 2.40 ± 0.33, and in patients with CT-SS ≥ 19.5 was 2.99 ± 0.47. Based on the independent T-test results between the intralesional vascular sizes of the two CT-SS categories, a statistically significant difference was found between the two (P = 0.001). In addition, the mean extralesional vascular size in patients with CT-SS < 19.5 was 2.04 ± 0.28, and in patients with CT-SS ≥19.5 was 2.35 ± 0.39. Based on the independent T-test results between the intralesional vascular sizes of the two CT-SS categories, a statistically significant difference was found between the two (P = 0.001).

**DISCUSSION**

This study showed a difference in vascular thickness between patients with mild and severe confirmed COVID-19 symptoms based on CT-SS scoring. It was found that distal blood vessel vascular thickening was more than average in intralesional and extralesional, and vascular thickening on CT-SS mild/severe intralesional was greater than extralesion. These results provide clinical implications that Covid 19 does not only involve respiratory organs, but systemic or multiorgan abnormalities occur. This result agrees with a previous study by Yang et al., who found an association between higher CT-SS scores and more severe COVID-19 infection. said that this feature of pulmonary vascular thickening is thought to contribute to respiratory failure in COVID-19 pneumonia. Vascular thickening is also associated with an increased prevalence of ARF, and the area of the lung parenchyma is affected. Several other investigators, Abbasi et al., found higher mortality rates in patients with higher CT-SS. The study by Provencher et al. showed that hospitalized COVID-19 patients presenting with pulmonary vascular thickening were the primary pulmonary abnormality in confirmed COVID-19 patients.<sup>6,8,7,9</sup>

On a normal chest CT scan, the subsegmental vessels distal to the lung should be inconspicuous in the subpleural region. According to Lang et al., pulmonary vascular thickening is defined as a larger diameter blood vessel characterized by: (a) the diameter of the blood vessel is more significant than that of the adjacent part of the lung that is not affected, (b) the diameter of the

vessel is more significant than that of the adjacent lung. In a comparable area of the unaffected contralateral lung, or (c) focal dilatation or non-tapping of blood vessels as they travel to the periphery of the lung. According to Tsujikawa et al., their study found the normal size value of the secondary pulmonary lobular vascular diameter to be +/- 0.5 - 1 mm. This study showed the pulmonary vascular wall thickening above the normal value.<sup>10,11</sup>

This vascular thickening is thought to be due to the role of endothelial cells and the mechanism of endothelial cell dysfunction in COVID-19. SARS-CoV-2 infects endothelial cells via angiotensin-converting enzyme 2 (ACE2)-mediated viral entry, facilitated by TMPRSS2, which triggers the SARS-CoV-2 spike glycoprotein. Endothelial cell infection can lead to downregulation of ACE2, leading to an imbalance between ACE2 and angiotensin II (AngII) levels, increasing Ang II. In addition, infection of endothelial cells or pericytes will interfere with crosstalk between these two cells, thereby contributing to endothelial cell dysfunction. In severe cases of COVID-19, activated macrophages release a variety of cytokines (e.g., soluble interleukin 2 receptors [IL-2R], interleukin-6 [IL-6], and Tumor Necrotizing Factor [TNFs]), which are associated with Cytokine Storm. It is mediated by the immune system and can cause vascular inflammation (endothelitis) due to increased adhesion molecules' expression on endothelial cells and inter-endothelial spaces, thereby increasing vascular hyperpermeability. This rapid spike in proinflammatory cytokines triggers inflammatory infiltration by lung tissue that causes lung damage to the epithelium and endothelium.<sup>8,12-15</sup>

In this study, it was found that there was a relationship between the appearance of pulmonary vascular thickening and the CT severity score in patients with confirmed COVID-19, where the appearance of vascular thickening increased with the presence of mild or severe CT-severity seen on the test.<sup>16-18</sup>

**CONCLUSION**

There is a relationship between pulmonary vascular thickening and CT Severity score on a CT scan of the thorax of patients with confirmed COVID-19. The more severe the CT Severity score in patients with confirmed COVID-19, the higher the findings of pulmonary vascular thickening on a chest CT scan.

## ACKNOWLEDGEMENT

Nothing to declare

## CONFLICT OF INTEREST

Nothing to declare

## SOURCE OF FUNDING

Nothing to declare

## REFERENCES

1. World Health Organization. Coronavirus disease (COVID-19). (2021).
2. Provencher, Steeve, François Potus, dan Sébastien Bonnet. 2020. COVID-19 and the pulmonary vasculature. *Pulmonary Circulation* 10.3: 2045894020933088.
3. Wang, Chuanbin, dkk. 2020. Initial CT features and dynamic evolution of early-stage patients with COVID-19. *Radiology of Infectious Diseases* 7.4: 195-203.
4. Zhao dkk 2020. Relation Between Chest CT Findings and Clinical Conditions of Coronavirus Disease (COVID-19) Pneumonia: A Multicenter Study
5. Zhou, Fei dkk. 2020. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The lancet* 395.10229: 1054- 1062.
6. Yang, Ran, dkk. 2020. Chest CT severity score: an imaging tool for assessing severe COVID-19. *Radiology: Cardiothoracic Imaging* 2.2: e200047.
7. Abbasi, Bitu dkk. Evaluation of the relationship between inpatient COVID-19 mortality and chest CT severity score. *The American Journal of Emergency Medicine*. 2021; 45: 458- 463.
8. Gonagle dennis mc, james s o'donnell, kassem sharif, paul emery, charles bridgewood immune mechanisms of pulmonary intravascular coagulopathy in covid-19 pneumonia . *lancet rheumatol* 2020 published online may 7, 2020
9. Provencher, Steeve, François Potus, dan Sébastien Bonnet. 2020. COVID-19 and the pulmonary vasculature. *Pulmonary Circulation* 10.3: 2045894020933088
10. Lang, Min dkk. 2020. Pulmonary vascular manifestations of COVID-19 pneumonia. *Radiology: Cardiothoracic Imaging* 2.3: e200277
11. Tsujikawa Tetsuya, dkk 2021, Radiological perspective of COVID-19 pneumonia: The early features and progressive behaviour on high-resolution CT. *Journal of Medical Imaging and Radiation Oncology* 65 208–212
12. Zhang, Haibo, dkk. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive care medicine* 2020. 46.4: 586-590.
13. Asakura Ogawa.2020 COVID-19-associated coagulopathy and disseminated intravascular coagulation. *International Journal of Hematology* volume 113
14. Guler, Nil, Fakiha Siddiqui, dan Jawed Fareed. 2020. Is the reason of increased D-dimer levels in COVID-19 because of ACE-2-induced apoptosis in endothelium. *Clinical and Applied Thrombosis/Hemostasis* 26: 1076029620935526
15. Lodigiani et al. Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res*. 2020;191:9- 14.
16. Mahardhika dkk, 2021. High D-dimer and CRP Levels in an Asymptomatic COVID-19 Patient: A Case Report and Brief Literature Review. *Seminar Nasional Riset Kedokteran 2 (SENSORIK)*
17. Wichmann et al. Autopsy findings and venous thromboembolism in patients with COVID-19. *Ann Intern Med*. 2020;173(4):268-277.
18. Yao, Yumeng dkk. 2020. D-dimer as a biomarker for disease severity and mortality in COVID-19 patients: a case control study. *Journal of intensive care* 8.1: 1-11.



